arXiv:1301.6349v3 [math.RA] 3 Apr 2013

Construction of Nilpotent Jordan Algebras Over any Arbitrary Fields

A.Hegazi , H.Abdelwahab
Mathematics department, Faculty of Science, Mansoura University, Egypt
hegazi@mans.edu.eg
hanii_i@yahoo.com

Abstract

We give a computional method to construct and classify nilpotent Jordan algebras over any arbitrary fields by the second cohomolgy of nilpotent Jordan algebras of low dimension "analogue of *Skjelbred-Sund method"* , we see that every nilpotent Jordan algebras can be constructed by the second cohomolgy of nilpotent Jordan algebras of low dimension. We use this method to classify nilpotent Jordan algebras up to dimension three over any field and nilpotent Jordan algebras of dimension four over an algebraic closed field of characteristic $\neq 2$ and over the real field $\mathbb{R}$. Also commutative nilpotent associative algebras "associative nilpotent Jordan algebras" are classified , we show that there are up to isomorphism 13 nilpotent Jordan algebras of dimension 4 over an algebraic closed field of characteristic $\neq 2$, and 4 of those are not associative, yielding 9 commutative nilpotent associative algebras. Also up to isomorphism there are 17 nilpotent Jordan algebras of dimension 4 over the real field $\mathbb{R}$, and 5 of those are not associative, yielding 12 commutative nilpotent associative algebras.

Keywords : Jordan algebras, Nilpotent, Centeral extension, Cohomology, Isomorphism, Automorphism group.

2010 Mathematics Subject Classification : 17C10, 17C55, 17-08.

1 Introduction

Jordan algebras were introduced in the early 1930's by a physicist, P. Jordan, in an attempt to generalize the formalism of quantum mechanics. Little appears to have resulted in this direction, but un-anticipated relationships between these algebras and Lie groups and the foundations of geometry have been discovered.

The structure of this paper, in section 2 we define the notion of Extension of Jordan algebra which is similar to the concept of Extension of Groups, let N, F and G be groups and let G have a normal subgroup $\bar{N}$ which is isomorphic to N, that is, $\bar{N} \cong N$. Recall that a subgroup is called normal or invariant if one has $g\bar{N}g^{-1} = \bar{N}$ for all $g \in G$. (At the Jordan algebras level a normal subgroup yields an ideal.) The group G is called an extension of F by N if the factor group G/N is isomorphic with the group F, i.e. $G/\bar{N} \cong F$. The relationship between the groups N, F and G can be represented by the sequence

$$N \xrightarrow{\Gamma} G \xrightarrow{\Lambda} F$$

where Γ is an injective group homomorphism with $\mathrm{im}\Gamma = \bar{N}$ and where Λ is a surjective group homomorphism with $\ker\Lambda = \bar{N}$. A systematic study of group extensions has been performed by the German mathematician O. Schreier in 1926.

In section 3, we focus on the Centeral Extension and in sections $4, 5$ and 6 we descripe the analouge of of *The Skjelbred-Sund Method,* and carry out a procudure for constructing nilpotent Jordan algebras over any arbitrary field. Finally, classify nilpotent Jordan algebras of dimension ≤ 4.

Definition 1 *A Jordan algebra J is a vector space over a field K equipped with a symmetric bilinear map $B : J \times J \longrightarrow J$ such that :*

$$B(B(x,x), B(x,y) = B(x, B(B(x,x),y)) \text{ , all } x, y \in J.$$

which is called the Jordan identity.

Let J be a Jordan algebra over a field K.For any integer $m \in \mathbb{N}$ we define a lower centeral series as the descending chain of ideals

$$c^1(J) = J \supseteq c^2(J) = B(J,J) \supseteq c^3(J) = B(c^2(J), J) \supseteq ..... \supseteq c^m(J) = B(c^{m-1}(J), J)$$

Definition 2 *A Jordan algebra J over a field K is said to be nilpotent of nilindex n if :*

$$c^n(J) = \{0\} \quad \textit{and} \quad c^{n-1}(J) \neq \{0\}.$$

Also we call n the nilpotency class of J and J is called n-step nilpotent.

Definition 3 *Let J be a Jordan algebra over a field K then the ideal :*

$$Z(J) = \{x \in J : B(x,y) = 0 \ \forall x \in J\}$$

is called the centre of J.

So, any nilpotent Jordan algebra have non-trivial centre $Z(J) \neq \{0\}$.In the Jordan algebra J, we will denote the product $B(x, y)$ by $x \circ y$ and x^2 for $B(x, x)$.Then the jordan identity is written as follow :

$$x^2 \circ (x \circ y) = x \circ (x^2 \circ y) \quad \text{, all } x, y \in J.$$

2 Extensions of Jordan algebras

In this section we are going to focus on extensions of Jordan algebras. Loosely speaking an extension of a Jordan algebra J is an enlargement of J by some other Jordan algebra. To be somewhat more concrete we consider the following construction. Starting with two Jordan algebras L and M over the same field K we consider the Cartesian product $J = L \times M$. Elements of this set are ordered pairs (l, m) with $l \in L$ and $m \in M$. Defining addition of such pairs by $(l, m) + (l', m') := (l + l', m + m')$ and multiplication by scalars $\alpha \in K$ as $\alpha(l, m) := (\alpha l, \alpha m)$ the set J becomes a vector space which will also be denoted by J. Using the Jordan multiplication on L and M we define on J the multiplication

$$(l, m) \circ (l', m') = (l \circ_L l', m \circ_M m')$$

One easily sees that this is a Jordan multipliction on J. Defining next the map $\epsilon : m \in M \longrightarrow (0, m) \in J$ and the map $\lambda : (l, m) \in J \longrightarrow m \in M$ one readily verifies that ϵ is an injective Jordan algebra homomorphism, while λ is a surjective Jordan algebra homomorphism. Moreover, im ϵ = ker λ. The Jordan algebra J with these properties is called a trivial extension of of L by M (or of M by L) .

Instead of denoting the elements of L by ordered pairs we will frequently use the notation $(l, m) = l + m$. The vector space J is then written as

$$J = L \oplus M$$

and the Jordan multiplication is in this notation given by

$$(l + m) \circ (l' + m') = l \circ_L l' + m \circ_M m'$$

This example of a trivial extension is generalized in the following concept. Let L, J and M be Jordan algebras and let these algebras be related in the following way.

- There exists an injective Jordan algebra homomorphism

$$\epsilon : L \longrightarrow J \ .$$

- There exists an surjective Jordan algebra homomorphism

$$\lambda : J \longrightarrow M \ .$$

- The Jordan algebra homomorphisms ϵ and λ are related by

$$\text{im } \epsilon = \text{ker } \lambda \ .$$

Then the Jordan algebra J is called an extension of M by L. The relationship is summarized by the sequence

$$L \xrightarrow{\epsilon} J \xrightarrow{\lambda} M$$

We elaborate a little on this concept. Since λ is a jordan algebra homomorphism one obtains that ker λ is an ideal in J and since λ is surjective First Isomorphism Theorem entails a Jordan algebra isomorphism between the quotient algebra $J/\text{ker } \lambda$ and L ,

$$J/\text{ker } \lambda \cong M$$

Using im ϵ = ker λ this relation can be written as

$$J/\text{im } \epsilon \cong M$$

Since ϵ is injective the Jordan algebras L and im ϵ are Jordan isomorphic, i.e $J/L \cong M$. From these properties one sees that it makes sense to call the Jordan algebra J is an extension of M by L.

Definition 4 *Let L, J and M be Jordan algebras over the field K. Let*

$$\lambda : J \longrightarrow M$$

be a surjective Jordan algebra homomorphism and

$$\epsilon : L \longrightarrow J$$

an injective Jordan algebra homomorphism. Then the sequence

$$L \xrightarrow{\epsilon} J \xrightarrow{\lambda} M \qquad (1)$$

is called an ***extension*** *of M by L if ϵ maps L onto the kernel ker $\lambda \subset J$ of the map λ. That is if*

$$im \ \epsilon = ker \ \lambda \ .$$

The kernel ker λ is called the ***kernel of the extension****.*

Loosely speaking J itself, instead of the sequence (1), will be called an extension of M by L.

It may happen that there exist several extensions of M by L. To classify extensions we define the notion of **equivalent extensions**.

Definition 5 *Two sequences*

$$L \xrightarrow{\epsilon} J \xrightarrow{\lambda} M$$

and

$$L \xrightarrow{\epsilon^{'}} J^{'} \xrightarrow{\lambda^{'}} M$$

are called ***equivalent extensions*** *if there exists a Jordan algebra isomorphism* $\phi : J \longrightarrow J^{'}$ *such that*

$$\phi \circ \epsilon = \epsilon^{'}, \quad \lambda^{'} \circ \phi = \lambda \ .$$

One easily sees that **equivalence of extensions** is an equivalence relation.

The concept of a Jordan algebra extension can be formulated more succinctly using the concept of an exact sequence.Let $\{J_i\}$ be a sequence of Jordan algebras and $\{\phi_i\}$ a sequence of Jordan algebra homomorphisms $\phi_i : J_i \longrightarrow J_{i+1}$, then the sequence

$$..... \longrightarrow J_{i-1} \xrightarrow{\phi_{i-1}} J_i \xrightarrow{\phi_i} J_{i+1} \xrightarrow{\phi_{i+1}} .....$$

is called **exact** if one has for each i

$$\text{im } \phi_{i-1} = \text{ker } \phi_i \ .$$

Let L, J and M be Jordan algebras and let us denote by 0 the Jordan algebra consisting of the zero element only. Then the sequence $0 \xrightarrow{f} L \xrightarrow{g} J$ is exact if and only if g is an injective Jordan algebra homomorphism. Indeed, since im $f = 0 \in L$ the requirement ker $g =$ im f forces g to be injective. One usually omits the mapping f i.e., one writes $0 \longrightarrow L \xrightarrow{g} J$. Likewise the sequence $J \xrightarrow{f} M \longrightarrow 0$ is exact if and only if f is a surjective Jordan algebra homomorphism. Consequently we have the following proposition.

Proposition 6 *The sequence* $L \longrightarrow J \longrightarrow M$ *of Jordan algebras is an* ***extension*** *of* M *by* L *if and only if the sequence*

$$0 \longrightarrow L \xrightarrow{\epsilon} J \xrightarrow{\lambda} M \longrightarrow 0$$

is ***exact .***

Next we introduce some particular types of extensions. For this we need the following concept from linear algebra. Two subspaces V_1 and V_2 of a vector space V are called complementary if V is the direct sum of V_1 and V_2.

Definition 7 *An extension*

$$L \xrightarrow{\epsilon} J \xrightarrow{\lambda} M$$

is called:
trivial *if there exists an ideal $I \subset J$ complementary to ker λ, i.e.*

$$J = ker\ \lambda \oplus I \qquad (\text{ Jordan algebra direct sum }),$$

split *if there exists a Jordan subalgebra $S \subset J$ complementary to ker λ, i.e.*

$$J = ker\ \lambda \oplus S \qquad (\text{ vector space direct sum }),$$

centeral *if the kernel ker λ is contained in the center $Z(J)$ of J, i.e.*

$$ker\ \lambda \subset Z(J)\ .$$

3 2-Cocycles on Jordan algebra

In this section we take a closer look at central extensions of Jordan algebras. The Jordan algebras in this section will be over the field K . Recall that the sequence of Jordan algebras

$$L \xrightarrow{\epsilon} J \xrightarrow{\lambda} M$$

is a central extension (of M by L) if one has for the Jordan algebra homomorphisms ϵ and λ the following properties: ϵ is injective, λ is surjective, im $\epsilon =$ ker λ and the kernel ker λ is contained in the center $Z(J)$ of J. Hence im $\epsilon =$ ker λ is a Jordan algebra with trivial multiplicatin. Since $Z(J)$ is a Jordan subalgebra with trivial multiplicatin and since L and im ϵ are isomorphic, L is a Jordan algebra with trivial multiplicatin too. A central extension of a Jordan algebra M by a Jordan algebra with trivial multiplicatin(a vector space) L can be obtained with the help of a so called 2-cocycle on M.

Definition 8 *Let J be a Jordan algebra and V be avector space over K. A bilinear map*

$$\theta : J \times J \longrightarrow V$$

is called a Jordan 2-cocycle from J to V if it satisfies for all $x, y \in J$ the following conditions :

- $\theta(x,y)=\theta(y,x)$ (θ is symmetric) .
- $\theta(x^2,x\circ y)=\theta(x,x^2\circ y)$ (Jordan identity for 2-cocycles) .

The set of 2-cocycles from J to V is denoted by $Z^2(J,V)$. One easily sees that $Z^2(J,V)$ is a vector space if one defines the vector space operations as follows. Let θ_1 and θ_2 be 2-cocycles rom J to V, then their linear combination $\lambda_1\ \theta_1+\ \lambda_2\ \theta_2$ ($\lambda_1,\lambda_2\in K$) is defined by

$$(\lambda_1\theta_1+\lambda_2\theta_2)(x,y):=\lambda_1\theta_1(x,y)+\lambda_2\theta_2(x,y)\ .$$

This linear combination is again a Jordan 2-cocycle.

Let $\theta\in Z^2(J,V)$,and set $J_\theta=J\oplus V$. Using Jordan multiplication on J and the cocycle θ from J to V e define on J_θ the multiplication

$$(x+v)\circ(y+w)=x\circ_J y+\theta(x,y)$$

For $x,y\in J,v,w\in V$.

Lemma 9 *J_θ is a Jordan algebra if and only if $\theta\in Z^2(J,V)$.*

Proof. For $x+v,y+w\in J_\theta$,we have that :

$$\begin{aligned}
(x+v)^2\circ((x+v)\circ(y+w)) &= (x\circ_J x+\theta(x,x))\circ(x\circ_J y+\theta(x,y))\\
&= x^2\circ_J(x\circ_J y)+\theta(x^2,x\circ_J y)\\
(x+v)\circ((x+v)^2\circ(y+w)) &= (x+v)\circ((x^2+\theta(x,x))\circ(y+w))\\
&= (x+v)\circ(x^2\circ_J y+\theta(x\circ_J x,y))\\
&= x\circ_J(x^2\circ_J y)+\theta(x,x^2\circ_J y)
\end{aligned}$$

From Jordan identity it follows that $(x+v)^2\circ((x+v)\circ(y+w))=(x+v)\circ((x+v)^2\circ(y+w)$ if and only if $\theta(x^2,x\circ y)=\theta(x,x^2\circ y)$ then J_θ is a Jordan algebra if and only if $\theta\in Z^2(J,V)$. ■

Lemma 10 *Let $\theta\in Z^2(J,V)$ then J_θ is a centeral extension of J by V .*

Proof. From the previous lemma J_θ is a Jordan algebra and V is a Jordan algebra with the trivial multiplication . The sequence $V\xrightarrow{i}J\oplus V\xrightarrow{\pi}J$ is exact(i be the injection map and π be the projection map) and $V=$ker $\pi\subset Z(J_\theta)$. Hence J_θ is a centeral extension of J by V . ■

Hence, a Jordan 2-cocycle θ from J to V (dim$V=k$) allows for the construction of a k-dimensional central extension $J_\theta=J\oplus V$ where the Jordan multiplication on J_θ is given by

$$(x+v)\circ(y+w)=x\circ_J y+\theta(x,y)$$

For $x, y \in J, v, w \in V$.

Now we will show that a Jordan 2-cocycles can be obtained from the"abstract form" of a central extension. Considering the extension

$$L \xrightarrow{\epsilon} J \xrightarrow{\lambda} M$$

with im ϵ =ker $\lambda \subset Z(J)$ we show that one can find 2-cocycles from M to L. Consider a linear map $s : M \longrightarrow J$ satisfying

$$\lambda \circ s = id_M$$

A map with this property is called a section of J. With the help of a section one can define a bilinear map $\theta' : M \times M \longrightarrow J$ by taking for all $x, y \in M$

$$\theta'(x, y) = s(x \circ y) - s(x) \circ s(y) \tag{2}$$

Notice that θ' is identically zero if s is a Jordan algebra homomorphism. Notice also that θ' is symmetric. From equation $\lambda \circ s = id_M$ and fact that λ is a Jordan algebra homomorphism one sees that

$$\lambda(\theta'(x, y)) = 0$$

Hence, we have for all $x, y \in M$

$$\theta'(x, y) \in \ker \lambda \subset Z(J) \ . \tag{3}$$

Using the injectivity of the map $\epsilon : L \longrightarrow J$ to define the map

$$\theta : M \times M \longrightarrow J \ .$$

Given by

$$\theta := \epsilon^{-1} \circ \theta'.$$

θ is bilinear and symmetric . It remains to show that θ satisfy the second condition in Defenition 8 . Let $x, y \in M$ then by using Jordan identity, equation (2) and property (3) we have

$$\begin{aligned}
\theta'(x^2, x \circ y) &= s(x^2 \circ (x \circ y)) - s(x^2) \circ s(x \circ y) \\
&= s(x \circ (x^2 \circ y)) - s(x^2) \circ (\theta'(x, y) + s(x) \circ s(y)) \\
&= s(x \circ (x^2 \circ y)) - s(x^2) \circ (s(x) \circ s(y)) \\
&= s(x \circ (x^2 \circ y)) - s(x) \circ (s(x^2) \circ s(y)) \\
&= s(x \circ (x^2 \circ y)) - s(x) \circ (\theta'(x^2, y) + s(x^2 \circ y)) \\
&= s(x \circ (x^2 \circ y)) - -s(x) \circ s(x^2 \circ y) \\
&= \theta'(x, x^2 \circ y) \ .
\end{aligned}$$

Hence, $\theta(x^2, x \circ y) = \epsilon^{-1}(\theta'(x^2, x \circ y)) = \epsilon^{-1}(\theta'(x, x^2 \circ y)) = \theta(x, x^2 \circ y)$. Then θ is a 2-cocycles from M to L. We have thus shown that 2-cocycles are quite natural objects in central extensions.

Corollary 11 *J_θ is a centeral extension of J by V if and only if $\theta \in Z^2(J,V)$.*

We now proceed with the properties of cocycles. A special type of 2-cocycles is given by so-called 2-coboundaries.

Definition 12 *Let J be a Jordan algebra and V be avector space over K. Then a linear map*

$$f : J \longrightarrow V.$$

is called a 1-cochain from J to V. The set of all 1-cochains from J to V is denoted by $C^1(J,V)$.

Notice that $C^1(J,V)$ be the vector space $Hom(J,V)$ and hence it has, just as$Z^2(J,V)$, an obvious vector space structure. Using cochains one defines 2-coboundaries as follows.

Definition 13 *Let J be a Jordan algebra,V be avector space over K and let $f : J \longrightarrow V$ be a 1-cochain from J to V. Then the bilinear map*

$$\delta f : J \times J \longrightarrow V.$$

defined by

$$(\delta f)(x,y) := f(x \circ y).$$

is called a $2-$coboundary from J to V.

Next we show that δf is a 2-cocycle from J to V. The symmetry of δf follows from

$$(\delta f)(x,y) := f(x \circ y) = f(y \circ x) = (\delta f)(y,x).$$

Furthermore the Jordan identity of J imply

$$(\delta f)(x^2, x \circ y) = f(x^2 \circ (x \circ y)) = f(x \circ (x^2 \circ y)) = (\delta f)(x, x^2 \circ y).$$

for all $x, y \in J$. Hence a 2-coboundary from J to V is a 2-cocycle from J to V. Consequently, the map $\delta : f \longrightarrow \delta f$ is a map from $C^1(J,V)$ to $Z^2(J,V)$.

Definition 14 *The map*

$$\delta : f \in C^1(J,V) \longrightarrow \delta f \in Z^2(J,V).$$

where δf is defined by $(\delta f)(x,y) := f(x \circ y)$ for all $x, y \in J$,is called the coboundary operator .

Using the notions of 2-cocycles and 1-cochains, more precisely the vector spaces $Z^2(J, V)$ and $C^1(J, V)$, one defines the second cohomology group of a Jordan algebra J by$V$.

Definition 15 *Let J be a Jordan algebra,V be avector space over K. Then the quotient vector space*

$$H^2(J, V) := Z^2(J, V) \diagup \delta C^1(J, V)$$

is called the second cohomology group of J by V.

Elements of $H^2(J, V)$ are equivalence classes of 2-cocycles and two 2-cocycles θ_1 and θ_2 are called equivalent cocycles if they differ by a 2-coboundary, i.e. if $\theta_1 = \theta_2 +\ \delta f$ for some $f \in C^1(J, V)$. Equivalent 2-cocycles are called cohomologous.

Lemma 16 *Let θ_1 and θ_2 be equivalent Jordan cocycles and let $J_{\theta_1}, J_{\theta 2}$ be respectively the centeral extensions constructed with these Jordan cocycles .Then the centeral extensions J_{θ_1} and $J_{\theta 2}$ are are equivalent extensions.*

Proof. According to the definition of equivalent Jordan cocycles we have $\theta_1 = \theta_2 + \delta f$ with $f \in C^1(J, V)$ a 1-cochain. Define $\sigma : J_{\theta_2} \longrightarrow J_{\theta_2 + \delta f}$ by $\sigma(x+v) = x+f(x)+v$. Let $x+v \in \ker \sigma$ then $\sigma(x+v) = x+f(x)+v = 0$,hence $x = v = 0$. Thus $\ker \sigma = \{0\}$, this shows that σ is an invertible linear transformation. Morever,for all $x + v, y + w \in J_{\theta_2}$

$$\begin{aligned}
\sigma((x + v) \circ_{J_{\theta_2}} (y + w)) &= \sigma(x \circ_J y + \theta_2(x, y)) \\
&= x \circ_J y + f(x \circ_J y) + \theta_2(x, y) \\
&= x \circ_J y + \delta f(x, y) + \theta_2(x, y) \\
&= x \circ_J y + (\theta_2 + \delta f)(x, y) \\
&= x \circ_J y + \theta_1(x, y) \\
&= (x + f(x) + v) \circ_{J_{\theta_1}} (y + f(y) + w) \\
&= \sigma(x + v) \circ_{J_{\theta_1}} \sigma(y + w).
\end{aligned}$$

Then J_{θ_1} and J_{θ_2} are isomorphic. Hence, cohomologous 2-cocycles yield equivalent (central) extensions. ■

Corollary 17 *A cohomology class $\theta \in H^2(J, V)$ defines a central extension of the Jordan algebra J by V which is unique up to equivalence.*

The cocycle $\theta = 0 \in H^2(J, V)$ gives a trivial central extension of a Jordan algebra J by V. Namely

$$J_0 = J \oplus V \qquad\qquad (\text{ direct sum of Jordan algebras })$$

and the Jordan multiplication is given by

$$(x + V) \circ_{J_0} (y + w) = x \circ_J y.$$

for all $x, y \in J$ and $v, w \in V$.

For any $f \in C^1(J, V)$ a 1-cochains, $\delta f = 0 + \delta f$. Hence, 2-cocycles which are obtained from 1-cochains are cohomologous with the trivial 2-cocycle $\theta = 0$ we obtain the following corollary.

Corollary 18 *A central extension defined by a* $2-$*coboundary is equivalent with a trivial central extension.*

Let V be m-dimensional vector space then $Z^2(J, V) = Z^2(J, K^m) = Z^2(J, K)^m$ and $H^2(J, V) = H^2(J, K^m) = H^2(J, K)^m$. So for any $\theta \in Z^2(J, V)$ we may write $\theta = (\theta_1, \theta_2, ..., \theta_m) \in Z^2(J, K)^m$. Let $e_1, e_2, ..., e_m$ be a basis of V then $\theta(x, y) = \sum\limits_{i=1}^{m} \theta_i(x, y) e_i$ where $\theta_i \in Z^2(J, K)$, and θ is a $2-$coboundary if and only if all $\theta_i \in \delta C^1(J, K)$.

Lemma 19 *dim* $\delta C^1(J, K) =$*dim* J^2

Proof. Let J be n-dimensional Jordan algebra with basis $< x_1, x_2, ..., x_n >$ and J^* generated by the dual basis $< x_1^*, x_2^*, ..., x_n^* >$ defined by $x_i^*(x_i) = 1$ and $x_i^*(x_j) = 0$ if $i \neq j$.Let $< x_r, x_{r+1}, ..., x_{r+s} >$ be a basis of J^2.Notice that $C^1(J, K)$ is the dual vector space of J. The coboundary operator

$$\delta : J^* \longrightarrow Z^2(J, K)$$

defined by

$$(\delta x_k^*)(x_i, x_j) = x_k^*(x_i \circ x_j)$$

for $x_k^* \in J^{\ *}$ and $x_i, x_j \in J$. Let $x_k \notin J^2$ then $\delta\ x_k^* = 0$, so $\delta C^1(J, K)$ is spanned by $< x_r^*, x_{r+1}^*, ..., x_{r+s}^* >$ which have a dimension equal to the dimension of J^2. ■

4 Analouge of The Skjelbred-Sund Method

Let J be a Jordan algebra with non-trivial centre $Z(J)$. We will show that such algebra are centeral extensions of smaller Jordan algebras .

Lemma 20 *Every n-dimensional Jordan algebra with non-trivial centre is a centeral extension of a lower dimensional Jordan algebra .*

Proof. Let J be $n-$dimensional Jordan algebra with non trivial centre$Z(J)$, then $J/Z(J)$ is a Jordan algebra of smaller dimension . Let $\pi : J \longrightarrow J/Z(J)$ be a projection map, choose an injective linear map $\eta : J/Z(J) \longrightarrow J$ such that $\pi(\eta(\bar{x})) = \bar{x}$ for all $\bar{x} \in J/Z(J)$. For $\bar{x}, \bar{y} \in J/Z(J)$ we have $\eta(\bar{x}) \circ \eta(\bar{y}) - \eta(\bar{x} \circ \bar{y}) \in Z(J)$. This define a symmetric bilinear map

$$\theta : J/Z(J) \times J/Z(J) \longrightarrow Z(J)$$

defined by $\theta(\bar{x}, \bar{y}) = \eta(\bar{x}) \circ \eta(\bar{y}) - \eta(\bar{x} \circ \bar{y})$. Then

$$\begin{aligned}
\theta(\bar{x}^2, \bar{x} \circ \bar{y}) &= \eta(\bar{x}^2) \circ \eta(\bar{x} \circ \bar{y}) - \eta(\bar{x}^2 \circ (\bar{x} \circ \bar{y})) \\
&= (\eta(\bar{x}) \circ \eta(\bar{x}) - \theta(\bar{x}, \bar{x})) \circ (\eta(\bar{x}) \circ \eta(\bar{y}) - \theta(\bar{x}, \bar{y})) - \eta(\bar{x} \circ (\bar{x}^2 \circ \bar{y}) \\
&= \eta(\bar{x}) \circ (\eta(\bar{x})^2 \circ \eta(\bar{y})) - \eta(\bar{x} \circ (\bar{x}^2 \circ \bar{y}) \\
&= \eta(\bar{x}) \circ (\eta(\bar{x}^2) \circ \eta(\bar{y})) - \eta(\bar{x} \circ (\bar{x}^2 \circ \bar{y}) \\
&= \theta(\bar{x}, \bar{x}^2 \circ \bar{y}).
\end{aligned}$$

Hence, θ is $2-$cocycle. It remains to show that $(J/Z(J))_\theta = J/Z(J) \oplus Z(J)$ is isomorphic to J. Let $x \in J$, then x can uniquely written as $x = \eta(\bar{y}) + z$, where $\bar{y} \in J/Z(J)$ and $z \in Z(J)$. Define $\phi : J \longrightarrow (J/Z(J))_\theta$ by $\phi(x) = \bar{y} + z$. Then ϕ is bijective and

$$\begin{aligned}
\phi(x_1 \circ_J x_2) &= \phi((\eta(\bar{y}_1) + z_1) \circ_J (\eta(\bar{y}_2) + z_2)) \\
&= \phi(\eta(\bar{y}_1) \circ_J \eta(\bar{y}_2)) \\
&= \phi(\eta(\bar{y}_1 \circ_{J/Z(J)} \bar{y}_2) + \theta(\bar{y}_1, \bar{y}_2)) \\
&= \bar{y}_1 \circ \bar{y}_2 + \theta(\bar{y}_1, \bar{y}_2) \\
&= (\bar{y}_1 + z_1) \circ_{(J/Z(J))_\theta} (\bar{y}_2 + z_2) \\
&= \phi(x_1) \circ_{{}_{(J/Z(J))_\theta}} \phi(x_2)
\end{aligned}$$

Then ϕ is an isomorhism. ■

So in particular, Every n-dimensional nilpotent Jordan algebras is a centeral extension of a lower dimensional nilpotent Jordan algebra .

Definition 21 *Let* $\theta \in Z^2(J,V)$ *then*

$$\theta^{\perp} = \{x \in J : \theta(x,y) = 0 \textit{ for all } y \in J\}$$

is called the radical of θ *.*

Let us now fix a basis $\{e_1, ..., e_r\}$ of V . A cocycle $\theta \in Z_2(J,V)$ such that $\theta(x,y) = \sum\limits_{i=1}^{r} \theta_i(x,y)e_i$ where $\theta_i \in Z^2(J,K)$. Then $\theta^{\perp} = \theta_1^{\perp} \cap \theta_2^{\perp} \cap .... \cap \theta_r^{\perp}$

Lemma 22 *Let* $\theta \in Z^2(J,V)$ *then* $Z(J_\theta) = (\theta^{\perp} \cap Z(J)) \oplus V$.

Proof. Let $x+v \in Z(J_\theta)$ then $(x+v) \circ_{J_\theta} (y+w) = 0$ for all $y+w \in J_\theta$.Then $x \circ_J y + \theta(x,y) = 0$ for all $y+w \in J_\theta$.It follows that $x \in \theta^{\perp} \cap Z(J)$ then $x+v \in (\theta^{\perp} \cap Z(J)) \oplus V$.On the other hand , suppose that $x+v \in (\theta^{\perp} \cap Z(J)) \oplus V$ then for all $y+w \in J_\theta$ we have $(x+v) \circ_{J_\theta} (y+w) = x \circ_J y + \theta(x,y) = 0$ then $x+v \in Z(J_\theta)$. Then $Z(J_\theta) = (\theta^{\perp} \cap Z(J)) \oplus V$. ■

A subspace W of $H^2(J,K)$ is said to be allowable if $\bigcap\limits_{\theta \in W} \theta^{\perp} \cap Z(J) = 0$.

Corollary 23 $\theta^{\perp} \cap Z(J) = 0$ *if and only if* $Z(J_\theta) = V$.

If $\theta^{\perp} \cap Z(J) \neq 0$ then J_θ can be obtaines as a centeral extension of another Jordan algebra $\tilde{J}$ by $(\theta^{\perp} \cap Z(J)) \oplus V$. Then to avoid constructing the same Jordan algebra as a centeral extension of different Jordan algebra we want to restrict to θ such that $\theta^{\perp} \cap Z(J) = 0$.

Let J be a Jordan over a field K . For each $\theta \in Z^2(J,V)$ and $\phi \in Aut(J)$,the automorphism group of J , we define $\phi\theta(x,y) = \theta(\phi x, \phi y)$ for any $x,y \in J$. So $Aut(J)$ acts on $Z^2(J,V)$, and $\phi\theta \in \delta C^1(J,V)$ if and only if $\theta \in \delta C^1(J,V)$. Then $Z^2(J,V)$ and $\delta C^1(J,V)$ are invariant under the action of $Aut(J)$. So $Aut(J)$ acts on $H^2(J,V)$.

Let $\theta_1, \theta_2 \in H^2(J,V)$ and $\theta_1^{\perp} \cap Z(J) = \theta_2^{\perp} \cap Z(J) = 0$, i.e $Z(J_{\theta_1}) = Z(J_{\theta_2}) = V$. Assume that J_{θ_1} and J_{θ_2} are isomorphic, let $\alpha : J_{\theta_1} \longrightarrow J_{\theta_2}$ be an isomorphism .Dividing with the common center V we obtain an automorphism $\alpha_0 : J \longrightarrow J$. We can realize α as a matrix to a suitable basis for $J \oplus V$ which assumed to contain a basis for J and a basis for V :

$$\alpha = \begin{pmatrix} \alpha_0 & 0 \\ \varphi & \psi \end{pmatrix}$$

Where $\alpha_0 \in Aut(J)$, $\psi = \alpha \mid_V \in Gl(v)$ and $\varphi \in Hom(J,V)$.

Now α preserves the Jordan products and writing $\circ_1$and $\circ_2$ for the Jordan products of J_{θ_1} and J_{θ_2} respectively, we have

$$\alpha((x+v) \circ_1 (y+w)) = \alpha(x+v) \circ_2 \alpha(y+w) \ \ ; \ \ x,y \in J \text{ and } v,w \in V.$$

Where

$$\begin{aligned}\alpha(x+v) &= \alpha_0(x)+\varphi(x)+\psi(v)\\ \alpha(y+w) &= \alpha_0(y)+\varphi(y)+\psi(w).\end{aligned}$$

Then

$$\begin{aligned}\alpha((x+v)\circ_1(y+w)) &= \alpha(x\circ y+\theta_1(x,y))\\ &= \alpha_0(x\circ y)+\varphi(x\circ y)+\psi(\theta_1(x,y)).\\ \alpha((x+v)\circ_2\alpha(y+w)) &= \alpha_0((x)+\varphi(x)+\psi(v))\circ_2(\alpha_0(y)+\varphi(y)+\psi(w))\\ &= \alpha_0(x)\circ\alpha_0(y)+\theta_2(\alpha_0(x),\alpha_0(y))\end{aligned}$$

This yields

$$\begin{aligned}\theta_2(\alpha_0(x),\alpha_0(y)) &= \varphi(x\circ y)+\psi(\theta_1(x,y))\ \ ,\ x,y\in J\\ \theta_2(\alpha_0(x),\alpha_0(y)) &= \psi(\theta_1(x,y)) \mod \delta C^1(J,V)\end{aligned}$$

In case of $\theta_1=\theta_2=\theta$ we obtain the following description of $Aut(J_\theta)$

Lemma 24 *Let J be a nilpotent Jordan algebra .Let $\theta\in H^2(J,V)$ and $\theta^\perp\cap Z(J)=0$. Then the automorphism group $Aut(J_\theta)$ of the extension algebra J_θ consists of all linear operators of the matrix form*

$$\alpha=\begin{pmatrix}\alpha_0 & 0\\ \varphi & \psi\end{pmatrix}$$

Where $\alpha_0\in Aut(J)$, $\psi=\alpha\mid_V\in Gl(V)$ and $\varphi\in Hom(J,V)$. Such that :

$$\theta(\alpha_0(x),\alpha_0(y))=\varphi(x\circ y)+\psi\theta(x,y)\ ,\forall\ x,y\in J.$$

Thus J_{θ_1} and J_{θ_2} are isomorphic if and only if there exist $\alpha_0\in Aut(J)$ and $\psi\in Gl(V)$ such that $\alpha_0\theta_2=\psi\theta_1 \mod \delta C^1(J,V)$, i.e $\alpha_0\theta_2$ and $\psi\theta_1$ are cohomologous. Let $\theta_1=(\theta_{11},\theta_{12},.....,\theta_{1s})\in Z^2(J,V)$, $\theta_2=(\theta_{21},\theta_{22},.....,\theta_{2s})\in Z^2(J,V)$ and $e_1,e_2,..........,e_m$ be a basis of V then $\theta_1(x,y)=\sum\limits_{i=1}^{s}\theta_{1i}(x,y)e_i$ and $\theta_2(x,y)=\sum\limits_{i=1}^{s}\theta_{2i}(x,y)e_i$ .Suppose that J_{θ_1} and J_{θ_2} are isomorphic , then $\sum\limits_{i=1}^{s}\alpha_0\theta_{2i}(x,y)e_i=\sum\limits_{i=1}^{s}\theta_{1i}(x,y)\psi(e_i)$. Let $\psi(e_i)=\sum\limits_{j=1}^{m}a_{ji}e_j$ then $\sum\limits_{i=1}^{s}\alpha_0\theta_{2i}(x,y)e_i=\sum\limits_{i=1}^{s}\sum\limits_{j=1}^{m}\theta_{1i}(x,y)a_{ji}e_j$ hence :

$$\alpha_0\theta_{2i}(x,y)=\sum_{i=1}^{s}\theta_{1i}(x,y)a_{ji}\quad \mod \delta C^1(J,V) \tag{4}$$

It follows that $\alpha_0\theta_{2i}$ span the same subspace of $H^2(J,K)$ as the θ_{1i} . We have proved

Lemma 25 *Let* $\theta = (\theta_1, \theta_2, ..., \theta_m)$ *and* $\eta = (\eta_1, \eta_2, ..., \eta_m) \in H^2(J,V)$ *and* $\theta^{\perp} \cap Z(J) = \eta^{\perp} \cap Z(J) = 0$ *. Then* J_θ *and* J_η *are isomorphic if and only if there exist* $\phi \in Aut(J)$ *such that* $\phi\theta_i$ *span the same subspace of* $H^2(J,K)$ *as the* η_i *.*

i.e Two allowable subspaces give isomorphic Jordan algebras if and only if they are in the same $Aut(J)$-orbit.

Let $J = I_1 \oplus I_2$ be the direct sum of two ideals. Suppose that I_2 is contained in the centre of J. Then I_2 is called a centeral component of J.

Jordan algebra $J = I_1 \oplus I_2$ with centeral component I_2 is a trivial centeral extension of I_1 by I_2. i.e Jordan algebras with centeral components are simply obtained by taking direct sums of Jordan algebras of smaller dimension with trivial Jordan algebras (Vector spaces). Therefore when constructing Jordan algebra with non trivial centre as centeral extension we want to avoid constructing those with centeral components. The folowing Lemma will help us to exclude Jordan algebras wit centeral components .

Lemma 26 *Let* $\theta(x,y) = \sum\limits_{i=1}^{r} \theta_i(x,y)e_i \in H^2(J,V)$ *and* $\theta^{\perp} \cap Z(J) = 0$ *. Then* J_θ *has a centeral component if and only if* $\theta_1, \theta_2, ..., \theta_r$ *are linearly dependent .*

Proof. Suppose that $\theta_1, \theta_2, ..., \theta_r$ are linearly dependent, then there exist a proper subset of $\theta_1, \theta_2, ..., \theta_r$ span the same as θ_i. So we may assume that that some of θ_i are zero , hence J_θ has a centeral component .

On the other hand, Let $\theta(x,y) = \sum\limits_{i=1}^{r=s+t} \theta_i(x,y)e_i \in H^2(J,V)$ and $\theta^{\perp} \cap Z(J) = 0$.Suppose that $J_\theta = J \oplus V$ has a centeral component B , and B is contained in V . Write $J_\theta = J \oplus W \oplus B$, then J_θ can be considered as a trivial centeral extension of $\tilde{J} = J \oplus W$ by B .The centre of J_θ is equal to $V = W \oplus B$ .Consider $\{e_1, e_2, ..., e_s\}$ and $\{e_{s+1}, e_{s+2}, ..., e_t\}$ are basis of W and B respectively. Also $\tilde{J} = J \oplus W$ can be considered as a centeral extension of J by$W$, then there exist a Jordan cocycle $\psi(x,y) = \sum\limits_{i=1}^{s} \psi_i(x,y)e_i$

such that $\psi^{\perp} \cap Z(J) = 0$ and $J_\psi = \tilde{J} = J \oplus W$. Then J_θ is a centeral extension of $J_\psi$by B with trivial Jordan cocycle,, then

$$\theta(x,y) = \sum_{i=1}^{r=s+t} \theta_i(x,y)e_i = \psi(x,y) = \sum_{i=1}^{s} \psi_i(x,y)e_i$$

Hence $\theta_1, \theta_2, ..., \theta_r$ are lineary dependent . ■

So to exclude centeral components, we must have θ_i to be linearly independent .

A Jordan algebra $\tilde{J}$ is said to be a descendant of the Jordan algebra J if $\tilde{J}/Z(\tilde{J}) \cong J$ and $Z(\tilde{J}) \leq' \tilde{J}^2$. If dim $Z(\tilde{J}) = r$ then $\tilde{J}$ is also referred to as a step$-r$ descendant. A descendant of a nilpotent Jordan algebra is nilpotent. Conversely, if $\tilde{J}$ is a finite-dimensional nilpotent Jordan algebra over a field K, then by Lemma (20) $\tilde{J}$ is either a descendant of a smaller-dimensional nilpotent Jordan algebra when $\tilde{J}$ has no centeral components, or $\tilde{J} = J \oplus Kx$ where J is an ideal of $\tilde{J}$ and Kx is a centeral component of $\tilde{J}$ (Kx is a 1-dimensional Vector space viewed as trivial Jordan algebra).

All our previous observations can be summarized as follows:

Theorem 27 *Let J be a Lie algebra, let Vbe a vector space with fixed basis $\{e_1, ..., e_r\}$ over a field K, and let θ, η be elements of $Z^2(J,V)$.*

1. *The Jordan algebra J_θ is a step-r descendant of J if and only if $\theta^{\perp} \cap Z(J) = 0$ and the image of the subspace $< \theta_1, ..., \theta_r >$ in $H^2(J,K)$ is r-dimensional.*
2. *Suppose that η is an other element of $Z^2(J,V)$ and that J_θ, J_η are descendants of J. Then $J_\theta \cong J_\eta$ if and only if images of the subspaces $< \theta_1, ..., \theta_r >and< \eta_1, ..., \eta_r >$ in $H^2(J,K)$ are in the same orbit under the action of $Aut(J)$.*

It follows that there is a one-to-one correspondence between the set of isomorphism types of step-r descendants of J and the Aut(J)-orbits on the r-dimensional allowable subspaces of $H^2(J,K)$. Hence the classification of n-dimensional nilpotent Jordan algebras requires that we determine these orbits for all nilpotent Jordan algebras of dimension at most $n-1$.

Let $G_r(H^2(J,K))$ be the Grassmanian of subspaces of dimension r in $H^2(J,K)$. There is anatural action of $Aut(J)$ on $G_r(H^2(J,K))$ defined by :

$$\phi < \theta_1, \theta_2, ..., \theta_r > = < \phi\theta_1, \phi\theta_2, ..., \phi\theta_r >$$

for $V = < \theta_1, \theta_2, ..., \theta_r > \in G_r(H^2(J,K))$ and $\phi \in Aut(J)$.

Note that if $\{\theta_1, \theta_2, ..., \theta_r\}$ is linear independent so is$\{\phi\theta_1, \phi\theta_2, ..., \phi\theta_r\}$.Define

$$U_r(J) = \{V = < \theta_1, \theta_2, ..., \theta_r > \in G_r(H^2(J,K)) : \theta_i^{\perp} \cap Z(J) = 0; i = 1, 2, ..., r\}.$$

Lemma 28 *$U_r(J)$ is stable under the action of $Aut(J)$.*

Proof. Let $\phi \in Aut(J)$ and $V = < \theta_1, \theta_2, ..., \theta_r > \in U_r(J)$. Let $x \in (\phi\theta_i)^{\perp} = \{y \in J : \phi\theta_i(y, J) = 0\}$ then $\phi(x) \in \theta_i^{\perp}$ hence $x \in \phi^{-1}\theta_i^{\perp} = \{\phi^{-1}(y) : y \in \theta_i^{\perp}\}$. Also let $\phi^{-1}(x) \in \phi^{-1}\theta_i^{\perp}$ then $\phi\theta_i(\phi^{-1}(x), J) = \theta_i(x, J) = 0$ hence $\phi^{-1}(x) \in (\phi\theta_i)^{\perp}$. Then $(\phi\theta_i)^{\perp} = \phi^{-1}\theta_i^{\perp}$ and $\phi^{-1}(Z(J)) = Z(J)$. Therefore $(\phi\theta_i)^{\perp} \cap Z(J) = \phi^{-1}(\theta_i^{\perp} \cap Z(J)) = 0$ then $\phi V \in U_r(J)$. ■

Let $U_r(J)\diagup Aut(J)$ be the set of $Aut(J)$-orbits of $U_r(J)$. Then there exists a canonical one-to-one correspondense from $U_r(J)\diagup Aut(J)$ onto the set of isomorphism classes of Jordan algebras without centeral components which are central extensions of J by V and have r-dimensional center where $r = \dim V$.

All our previous observations can be summarized to have an analogue of the *Skejelbred-Sund theorem* for Jordan algebras as follows:

Theorem 29 *Let J be a Jordan algebra over a field K.The isomorphism clases of Jordan algebras $\tilde{J}$ with centre V of dimension r ,$\tilde{J}\diagup V \cong J$ and without centeral component are in bijective correspondence with the elements in $U_r(J)\diagup Aut(J)$.*

By this theorem, we may construct all nilpotent Jordan algebras of dimension n, given those algebras of dimension less than n, by centeral extension.

5 Constructing nilpotent Jordan algebras

In this section we carry out a procudure for constructing nilpotent Jordan algebras over any arbitrary field.

The classification procedure :

Let J be a Jordan algebras with basis $e_1, e_2, ..., e_{n-r}$ then the dual basis of the space of all symmetric bilinear forms $\theta : J \times J \longrightarrow K$ are $\sum_{e_i,e_j}$ $: n-r \geq i \geq j = 1$ with $\sum_{e_i,e_j}(e_l, e_m) = \sum_{e_i,e_j}(e_m, e_l) = 1$ if $i = l$ and $j = m$,and takes the value 0 otherwise, then any $2-$cocycle $\theta \in Z^2(J, K)$ can be represented by $\theta = \sum\limits_{i \geq j=1}^{n-r} c_{ij} \sum_{e_i,e_j}$ such that the elements $c_{i,j}$ satisfy Jordan identity for $2-$cocycles . By Theorem (29) we have a procdure that takes as input a Jordan algebra J of dimension $n-r$ it outputs all nilpotent Jordan algebras $\tilde{J}$ of dimension n such that $\tilde{J} \diagup Z(\ \tilde{J}) \cong J$, and $\tilde{J}$ has no centeral components. It runs as follows :

- For a given nilpotent Jordan algebra J of dimension $n-r$, we list at first its center to help us identify the $2-$cocycles satisfying $\theta^{\perp} \cap Z(J)$ $= 0$.
- Compute $Z^2(J, K)$: When computing the $2-$cocycles, we will just list all the constraints on the elements c_{ij}.
- Compute $\delta C^1(J, K)$: For $e_l \in J^2$ define $\theta_l = \sum\limits_{i,j} \left(\lambda^l_{i,j} \sum_{e_i,e_j} \right)$ such that $e_i \circ e_j = e_l$. Then $\delta C^1(J, K)$ is spanned by θ_l for all $e_l \in J^2$, the scalers $\lambda^l_{i,j}$ which at least one of them is nozero are determined by considering the invarient of $\delta C^1(J, K)$ under the action of the automorphism group.
- Compute $H^2(J, K)$: The complement of $\delta C^1(J, K)$ in $Z^2(J, K)$ be $H^2(J, K)$. For $\theta = \sum\limits_{i \geq j=1}^{n-r} c_{i,j} \sum_{e_i,e_j} \in Z^2(J, K)$ put $c_{i,j} = 0$ for all $\sum_{e_i,e_j} = \theta_l \in \delta C^1(J, K)$.
- Consider $\theta \in H^2(J, V)$ with $\theta(x, y) = \sum\limits_{i=1}^{r} \theta_i(x, y) e_{n-r+i}$ where the $\theta_i \in H^2(J, K)$ are linearly indpendent, and $\theta^{\perp} \cap Z(J) = 0$. Find a list of representatives of the orbits of $Aut(J)$ acting on the θ.

- For each θ found, construct J_θ. Discard the isomorphic ones.

Remarks :

1. If charcteristic $K \neq 2$, one can linearize $\theta(x^2, x \circ y) = \theta(x, x^2 \circ y)$ to obtain :

   $$\theta(x, v\circ(y\circ z))+\theta(y, v\circ(x\circ z))+\theta(z, v\circ(x\circ y)) = \theta(x\circ y, z\circ v)+\theta(y\circ z, x\circ v)+\theta(x\circ z, y\circ v)$$

   for all $x, y, z, v \in J$. May be usefel for computing $Z^2(J, K)$.

2. Let $J : a^2 = c, b^2 = -c$, choosing one of $\sum_{a,a}$ or $\sum_{b,b}$ to be a basis for the coboundary space,$\delta C^1(J, K)$, is not correct. Because if we choose the coboundary space spanned by $\sum_{b,b}$, we will get the following algebras

   $$\begin{aligned} J_{\theta_1} &: a^2 = c, b^2 = -c, a \circ c = d, b \circ c = d \\ J_{\theta_2} &: a^2 = c, b^2 = -c - d, a \circ c = d, b \circ c = d \end{aligned}$$

   isomorphic since $\theta_1 - \theta_2 = \sum_{b,b}$ which is not true. Also if we consider the coboundary space,$\delta C^1(J, K)$, spanned by $\sum_{a,a}$ and $\sum_{b,b}$ is also not correct since dim $\delta C^1(J, K)$ =dim J^2.

3. One can use Lemma (24) to compute the automorphism group,and the system of equations (4)for testing if two cocycles are in the same $Aut(J)$-orbit by checking solvalbality. However in the next section we construct an algorithm for testing isomorphism of any two Jordan algebra, by this algorithm we can also compute the automorphism group.

4. The procedure only gives those nilpotent Jordan algebras without cetral components. So we have to add the nilpotent Jordan algebras obtained by by taking direct sum of a smaller dimensional nilpotent Jordan algebra with trivial Jordan algebra (that has trivial multiplication).

6 Deciding isomorphism of Jordan algebras

A classical problem is to know how many different (up to isomorphisms) finite dimensional Jordan algebras exist for each dimension. Let J be a

Jordan algebra of dimension n, then it has a basis $e_1, e_2, ..., e_n$. It follows that there are constants c_{ij}^k for $n \geq i \geq j = 1$ such that

$$e_i \circ e_j = \sum_{k=1}^{n} c_{ij}^k e_k.$$

Let J_1 and J_2 be two Jordan algebras for which we want to decide whether they are isomorphic or not. If the dimension of J_1 and J_2 are not equal, then J_1 and J_2 can never be isomorphic. Furthermore, suppose that $\phi : J_1 \longrightarrow J_2$ is an isomorphism of Jordan algebras, then $\phi(Z(J_1)) = Z(J_2)$ and $\phi(C^k(J_1)) = C^k(J_2)$. So if $\dim Z(J_1) \neq \dim Z(J_2)$ then J_1 and J_2 can never be isomorphic. Also if $\dim C^k(J_1) \neq \dim C^k(J_2)$ hen J_1 and J_2 can never be isomorphic. By this way we may able to decide that J_1 and J_2 are not isomorphic. There is a direct method for testing isomorphism of Jordan algebras by using Grobner bases. Let $e_1, e_2, ..., e_n$ be a basis of J_1 and let $\tilde{e}_1, \tilde{e}_2, ..., \tilde{e}_n$ be a basis of J_2. Let (c_{ij}^k) and (γ_{ij}^k) be the structure constants of J_1 and J_2, respectively. A map $\phi : J_1 \longrightarrow J_2$ is an isomorphism of Jordan algebras if and only if it satisfies the following requirements :

- $\phi(e_i \circ e_j) = \phi(e_i) \circ \phi(e_j)$ for $n \geq i \geq j = 1$,
- ϕ is non-singular.

Let $\phi : J_1 \longrightarrow J_2$ is an isomorphism of Jordan algebras given by $\phi(e_i) = \sum_{j=1}^{n} a_{ij}\tilde{e}_j$ then

$$\phi(e_i \circ e_j) = \phi\left(\sum_{k=1}^{n} c_{ij}^k e_k\right) = \sum_{k=1}^{n} c_{ij}^k \phi(e_k) = \sum_{k,m=1}^{n} c_{ij}^k a_{km}\tilde{e}_m,$$

and

$$\begin{aligned} \phi(e_i) \circ \phi(e_j) &= \left(\sum_{k=1}^{n} a_{ik}\tilde{e}_k\right) \circ \left(\sum_{l=1}^{n} a_{jl}\tilde{e}_l\right) \\ &= \sum_{k,l=1}^{n} a_{ik}a_{jl}(\tilde{e}_k \circ \tilde{e}_l) = \sum_{k,l,m=1}^{n} \gamma_{kl}^m a_{ik}a_{jl}\tilde{e}_m. \end{aligned}$$

Hence the first requirement amounts to the following $\frac{n(n+1)}{2}$ equations in the variables a_{ij} :

$$\sum_{k=1}^{n} c_{ij}^k a_{km} - \sum_{k,l=1}^{n} \gamma_{kl}^m a_{ik}a_{jl} = 0 \text{ for } n \geq i \geq j = 1 \text{ and } 1 \leq m \leq n.$$

By the second erquirement the determinant of the matrix (a_{ij}) not equal to 0, $\det(a_{ij}) \neq 0$.

Theorem 30 ***(Generalized Weak Nullstellensatz)****. Let k be any field. A system of polynomials $f_1, ..., f_m \in k[x1, ..., xn]$has no common zero over the algebraic closure of k, if and only if, $1 \in I$ where I be the ideal generated by the polynomials $f_1, ..., f_m$.*

So we consider the ideal I generated by the polynomials

$$\sum_{k=1}^{n} c_{ij}^{k} a_{km} - \sum_{k,l=1}^{n} \gamma_{kl}^{m} a_{ik} a_{jl} = 0 \text{ for } n \geq i \geq j = 1 \text{ and } 1 \leq m \leq n.$$

together with the polynomial $b \det(a_{ij}) - 1$ in the polynomial ring $k[a_{ij}, b]$. Then there is no solution of this system if and only if $I \neq k[a_{ij}, b]$. The algorithm for calculating Grobener bases yields a method for deciding whether or not $1 \in I$. Then by Grobener bases calculations we can decide isomorphism of Jordan algebras.

Algorithm 31 ***Testing isomorphism for Jordan algebras****.*

Input : Two Jordan algebras $(J_1, \circ_1)$ with basis $e_1, e_2, ..., e_n$ and $(J_2, \circ_2)$ with basis $\tilde{e}_1, \tilde{e}_2, ..., \tilde{e}_n$.

Output ; True if $J_1 \cong J_2$ and false in other case.

1. *Compute the following system of equations :*

   $$\phi(e_i \circ_1 e_j) - \phi(e_i) \circ_2 \phi(e_j) = 0 \textit{ for } n \geq i \geq j = 1.$$

   where $\phi = (a_{ij})_{n \times n}$

2. *To ensure that ϕ is going to be non-singular, add the following relation with a new variable b :*

   $$b \det \phi - 1 = 0.$$

3. *Compute a grobener bases G of the ideal $I =< \{\phi(e_i \circ_1 e_j) - \phi(e_i) \circ_2 \phi(e_j)\}_{n \geq i \geq j = 1} \cup \{b \det \phi - 1\} >$ in the polynomial ring $k[a_{ij}, b]$.*

4. *$G = \{1\}$?*

   (a) *Yes.*
   Return False.

(b) No.
Output : True ; Return G.

If the output of the algorithm is true, then it provides the equations of an algebraic variety whose points are all possiple values for ϕ, hence we have a descripiton of the automorphism group in the case $J_1 = J_2$.

7 Nilpotent Jordan algebras of dimension≤ 4

In this section we use the analouge **Skjelbred-Sund method** to classify nilpotent Jordan algebras of dimension ≤ 3 over any field, and four dimensional nilpotent Jordan algebras in the following cases :

- Over algebraic closed field K and $ch(K) \neq 2$.
- Over the real field $\mathbb{R}$.

Here we denote the j-th algebra of dimension i by $J_{i,j}$.

7.1 Nilpotent Jordan algebras of dimension 1

Let J be a Jordan algebra of dimension 1 spanned by a. If J is nilpotent then $Z(J)$ is non-trivial. Hence $Z(J)$ is spanned by a. It follows that there is only one nilpotent Jordan algebra of dimension 1 over any arbitrary field,$J_{1,1}$, it spanned by a and $a^2 = 0$.

$J_{1,1}$	All multiplications are zero.

7.2 Nilpotent Jordan algebras of dimension 2

To construct all nilpotent Jordan algebras of dimension 2, firstly we consider the trivial 1-dimensional Central extension of the nilpotent Jordan algebras of dimension 1 corresponding to $\theta = 0$ (Algebras that are direct sum of an algebra of dimension 1 and a 1-dimensional trivial Jordan algebra, isomorphic to $J_{1,1}$) to get nilpotent Jordan algebras with *centeral component* . Finally consider non trivial 1- dimensional Central extension of the nilpotent Jordan algebras of dimension 1 to get nilpotent Jordan algebras without *centeral component.*

7.2.1 Nilpotent Jordan algebras with Centeral component

1. **1- dimensional Trivial Central extension of $J_{1,1}$ corresponding to $\theta = 0$.**

   (a) We get the Jordan algebra :

$J_{2,1} = J_{1,1} \oplus J_{1,1}$	All multiplications are zero.

7.2.2 Nilpotent Jordan algebras without Centeral component

1. 1**- dimensional Central extension of** $J_{1,1}$

   Here we get that $Z^2(J_{1,1}, K)$ is spanned by $\sum_{a,a}$ and $\delta C^1(J_{1,1}, K) = 0$, then $H^2(J_{1,1}, K)$ is spanned by $\sum_{a,a}$. So we get only one cocycle $\theta = \sum_{a,a}$, yielding the algebra :

$J_{2,2}$	$a^2 = b.$

Theorem 32 *Up to isomorphism there exactly two nilpotent Jordan algebras of dimension* 2 *over any field* K *which are isomorphic to the following nilpotent Jordan algebras :*

Nilpotent Jordan algebras with Centeral component
$J_{2,1} = J_{1,1} \oplus J_{1,1}$.
Nilpotent Jordan algebras without Centeral component
$J_{2,2} : a^2 = b.$

7.3 Nilpotent Jordan algebras of dimension 3

From the previous subsection we get that the 2-dimensional nilpotent Jordan algebras over any field K are : $J_{2,1}$ and $J_{2,2}$. Using those algebra to construct 3-dimensional nilpotent Jordan algebras as follow :

7.3.1 Nilpotent Jordan algebras with Centeral component

1. **1- dimensional Trivial Central extension of $J_{2,1}$ corresponding to $\theta = 0$.**

   We get the Jordan algebra :

$J_{3,1} = J_{2,1} \oplus J_{1,1}$	All multiplications are zero.

2. 1**- dimensional Trivial Central extension of $J_{2,2}$ corresponding to $\theta = 0$.**

We get the Jordan algebra :

$J_{3,2} = J_{2,2} \oplus J_{1,1}$	$a^2 = b.$

7.3.2 Nilpotent Jordan algebras without Centeral component

1. 1- **dimensional Central extension of** $J_{2,1}$

   Here we get that $Z^2(J_{2,1}, K)$ is spanned by $\sum_{a,a}$,$\sum_{a,b}$ and $\sum_{b,b}$, also we get that $\delta C^1(J_{2,1}, K) = 0$. Then $H^2(J_{2,1}, K)$ is consists of $\theta = \alpha_1 \sum_{a,a} + \alpha_2 \sum_{a,b} + \alpha_3 \sum_{b,b}$. The centre of $J_{2,1}, Z(J_{2,1})$, spanned by a and b, it follows that $\theta : J_{2,1} \times J_{2,1} \longrightarrow K$ such that

   $$\theta^{\perp} \cap Z(J_{2,1}) = 0$$

   is nondegenerate since $\theta^{\perp} = 0$. The autommorphism group consists of :

   $$\phi = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}, \det \phi \neq 0.$$

   Characteristic $K \neq 2$

   Every symmetric bilinear form is diagonalizable, so we may assume that $\theta = \sum_{a,a} + \alpha \sum_{b,b}$ and $\alpha \neq 0$. Write $\phi\theta = \alpha_1' \sum_{a,a} + \alpha_2' \sum_{a,b} + \beta \sum_{b,b}$, then

   $$\begin{aligned} \alpha_1' &= a_{11}^2 + \alpha a_{21}^2 \\ \alpha_2' &= a_{11}a_{12} + \alpha a_{21}a_{22} \\ \beta &= a_{12}^2 + \alpha a_{22}^2 \end{aligned}$$

   To fix $\alpha_1' = 1$ and $\alpha_2' = 0$, choose $a_{21} = a_{12} = 0$ and $a_{11} = 1$. Then $\phi\left(\sum_{a,a} + \alpha \sum_{b,b}\right) = \sum_{a,a} + \alpha a_{22}^2 \sum_{b,b}$. This show that for any $\alpha, \beta, a_{22} \in K^*$ such that $\beta = \alpha a_{22}^2$, then $\theta_{1,\alpha}$ and $\theta_{1,\beta}$ are in the same $Aut(J_{2,1})$-orbit. On the other hand , suppose that $\theta_{1,\alpha}$ and $\theta_{1,\beta}$ are in the same $Aut(J_{2,1})$-orbit. Then there exist $\phi \in Aut(J_{2,1})$ and $\lambda \in K^*$ such that $\phi\theta_{1,\alpha} = \lambda\theta_{1,\beta}$, so we have

   $$\phi^t \begin{pmatrix} 1 & 0 \\ 0 & \alpha \end{pmatrix} \phi = \lambda \begin{pmatrix} 1 & 0 \\ 0 & \beta \end{pmatrix}$$

   it follows that $\beta = \left(\frac{\det \phi}{\lambda}\right)^2 \alpha$. Hence $\theta_{1,\alpha}$ and $\theta_{1,\beta}$ are in the same $Aut(J_{2,1})$-orbit if and only if there is a $\delta \in K^*$ with $\beta = \delta^2\alpha$. So we get the algebra

$J_{3,3}^{\alpha}$	$a^2 = c \text{ , } b^2 = \alpha c \text{ , } \alpha \in K^* \diagup (K^*)^2 .$

Over algebraic closed field K, $K^* \diagup (K^*)^2 = \{1\}$. So we get only one

$J_{3,3}$	$a^2 = c\ ,\ b^2 = c.$

Over real field $\mathbb{R}$, $\mathbb{R}^* \diagup (\mathbb{R}^*)^2 = \{\pm 1\}$. So we get two algebras

$J_{3,3}$	$a^2 = c\ ,\ b^2 = c.$
$J_{3,3}^{-1}$	$a^2 = c\ ,\ b^2 = -c.$

Over finite field $K, K^* \diagup (K^*)^2 = \{\pm 1\}$. So we get two algebras

$J_{3,3}$	$a^2 = c\ ,\ b^2 = c.$
$J_{3,3}^{-1}$	$a^2 = c\ ,\ b^2 = -c.$

Over rational field $\mathbb{Q}, \mathbb{Q}^* \diagup (\mathbb{Q}^*)^2$ is infinite. Then there is an infinite number of them.

Characteristic $K = 2$

If θ is alternate we obtain only one cocycle $\theta_1 = \sum_{a,b}$. On the other hand if θ is not alternate, then it is diagonalizable. In this case $K^* \diagup (K^*)^2 = \{1\}$. so we obtain only one cocycle $\theta_2 = \sum_{a,a} + \sum_{b,b}$. Clearly θ_1, θ_2 are inequivalent bilinear form, hence they not lie in the same $Aut(J_{2,1})$-orbit,[$\phi\theta_1 = (\det\phi)\theta_1$. i.e $\phi\theta_1$ is a multiple of θ_1,hence it is not conjugate to θ_2. Then$(J_{2,1})_{\theta_1}$ and $(J_{2,1})_{\theta_2}$ are not isomorphic]. We get two algebras

$J_{3,3}$	$a^2 = c\ ,\ b^2 = c.$
$J_{3,4}$	$a \circ b = c.$

2. **1- dimensional Central extension of** $J_{2,2}$

   Here we get that $Z^2(J_{2,1}, K)$ is spanned by $\sum_{a,a}$ and $\sum_{a,b}$. Moreover $\delta C^1(J_{1,1}, K)$ is spanned by $\sum_{a,a}$, then $H^2(J_{1,1}, K)$ is spanned by $\sum_{a,b}$. So we get only one cocycle $\theta = \sum_{a,b}$, yielding the algebra

$J_{3,4}$	$a^2 = b\ ,\ a \circ b = c.$

3. **2- dimensional Central extension of** $J_{1,1}$

   We have that $H^2(J_{1,1}, K)$ is 1-dimensional spanned by $\sum_{a,a}$, then there is no 2- dimensional Central extension of $J_{1,1}$.

We can summarize some of our results in the following theorems :

Theorem 33 *Up to isomorphism there exist* 4 *nilpotent Jordan algebras of dimension* 3 *over algebraic closed field* K *and* $ch(K) \neq 2$ *which are isomorphic to one of the following pairwise non-isomorphic nilpotent Jordan algebras :*

Nilpotent Jordan algebras with Centeral component
$J_{3,1} = J_{2,1} \oplus J_{1,1}$.
$J_{3,2} = J_{2,2} \oplus J_{1,1}$.
Nilpotent Jordan algebras without Centeral component
$J_{3,3} : a^2 = c \ , \ b^2 = c.$
$J_{3,4} : a^2 = b \ , \ a \circ b = c.$

Theorem 34 *Up to isomorphism there exist* 5 *nilpotent Jordan algebras of dimension* 3 *over field* K *and* $ch(K) = 2$ *which are isomorphic to one of the following pairwise non-isomorphic nilpotent Jordan algebras :*

Nilpotent Jordan algebras with Centeral component
$J_{3,1} = J_{2,1} \oplus J_{1,1}$.
$J_{3,2} = J_{2,2} \oplus J_{1,1}$.
Nilpotent Jordan algebras without Centeral component
$J_{3,3} : a \circ b = c.$
$J_{3,4} : a^2 = c \ , \ b^2 = c.$
$J_{3,5} : a^2 = b \ , \ a \circ b = c.$

Theorem 35 *Up to isomorphism there exist* 5 *nilpotent Jordan algebras of dimension* 3 *over* $\mathbb{R}$ *which are isomorphic to one of the following pairwise non-isomorphic nilpotent Jordan algebras :*

Nilpotent Jordan algebras with Centeral component
$J_{3,1} = J_{2,1} \oplus J_{1,1}$.
$J_{3,2} = J_{2,2} \oplus J_{1,1}$.
Nilpotent Jordan algebras without Centeral component
$J_{3,3}^{\alpha=\pm 1} : a^2 = c \ , \ b^2 = \alpha c. \qquad (\alpha = \pm 1)$
$J_{3,4} : a^2 = b \ , \ a \circ b = c.$

We are showed that also the number of nilpotent Jordan algebras of dimension 3 over finite field is five for any characteristic.

Theorem 36 *Every nilpotent Jordan algebra of dimension* ≤ 3 *is a commutative nilpotent associative algebra.*

It follows that all commutative nilpotent associative algebras of dimension ≤ 3 are classified, and all previous theorems are valid for commutative nilpotent associative algebras.

7.4 Nilpotent Jordan algebras of dimension 4

In this subsection we construct all nonisomorphic nilpotent algebras of dimension four :

- Over algebraic closed field K and characteristic $K \neq 2$.
- Over real field $\mathbb{R}$.

Here we denote the j-th algebra of dimension i over K and $\mathbb{R}$ by $J_{i,j}(K)$ and $J_{i,j}(\mathbb{R})$ respectively.

7.4.1 Nilpotent Jordan algebras with Centeral component

1. **1- dimensional Trivial Central extension of $J_{3,1}$ corresponding to $\theta = 0$.**

   We get the Jordan algebra :

$J_{4,1}(K) = J_{3,1} \oplus J_{1,1}$	All multiplications are zero.

$J_{4,1}(\mathbb{R}) = J_{3,1} \oplus J_{1,1}$	All multiplications are zero.

2. **1- dimensional Trivial Central extension of $J_{3,2}$ corresponding to $\theta = 0$.**

   We get the Jordan algebra :

$J_{4,2}(K) = J_{3,2} \oplus J_{1,1}$	$a^2 = b.$

$J_{4,2}(\mathbb{R}) = J_{3,2} \oplus J_{1,1}$	$a^2 = b.$

3. **1- dimensional Trivial Central extension of $J_{3,3}^{\alpha}$,$\alpha \in K^* \diagup (K^*)^2$,corresponding to $\theta = 0$.**

   We get the Jordan algebra :

$J_{4,3}(K) = J_{3,3} \oplus J_{1,1}$	$a^2 = c$, $b^2 = c$.

$J_{4,3}^{\alpha=\pm1}(\mathbb{R}) = J_{3,3}^{\alpha=\pm1} \oplus J_{1,1}$	$a^2 = c$, $b^2 = \alpha c$. $(\alpha = \pm 1)$

4. **1- dimensional Trivial Central extension of $J_{3,4}$ corresponding to $\theta = 0$.**

   We get the Jordan algebra :

$J_{4,4}(K) = J_{3,4} \oplus J_{1,1}$	$a^2 = b$, $a \circ b = c.$

$J_{4,4}(\mathbb{R}) = J_{3,4} \oplus J_{1,1}$	$a^2 = b$, $a \circ b = c.$

7.4.2 Nilpotent Jordan algebras without Centeral component

1. **1- dimensional Central extension of** $J_{3,1}$

   Here we get that $Z^2(J_{3,1},K)$ is spanned by $\sum_{a,a}, \sum_{b,b}, \sum_{c,c}, \sum_{a,b}, \sum_{a,c}$ and $\sum_{b,c}$, also we get that $\delta C^1(J_{3,1},K) = 0$. Then $H^2(J_{3,1},K)$ consists of $\theta = \alpha_1 \sum_{a,a} + \alpha_2 \sum_{b,b} + \alpha_3 \sum_{c,c} + \alpha_4 \sum_{a,b} + \alpha_5 \sum_{a,c} + \alpha_6 \sum_{b,c}$. The centre of $J_{3,1}, Z(J_{3,1})$, spanned by a, b and c, it follows that $\theta : J_{3,1} \times J_{3,1} \longrightarrow K$ such that

   $$\theta^{\perp} \cap Z(J_{3,1}) = 0$$

   is nondegenerate since $\theta^{\perp} = 0$. The autommorphism group,$Aut(J_{3,1})$, consists of :

   $$\phi = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix}, \det \phi \neq 0.$$

   Over algebraic closed field K and characteristic $K \neq 2$:

   Up to equivalence there is only one nondegenerate symmetric bilinear form, $\sum_{a,a} + \sum_{b,b} + \sum_{c,c}$. So there is only one $Aut(J_{3,1})$-orbit $\theta = \sum_{a,a} + \sum_{b,b} + \sum_{c,c}$, yielding the algebra

$J_{4,5}(K)$	$a^2 = d\ ,\ b^2 = d\ , c^2 = d.$

   Over real field $\mathbb{R}$:

   By *Sylvester's law of inertia and Signature,* up to equivalence there are four nondegenerate symmetric bilinear form

   $\theta_1 = \sum_{a,a} + \sum_{b,b} + \sum_{c,c}$
   $\theta_2 = \sum_{a,a} + \sum_{b,b} - \sum_{c,c}$
   $\theta_3 = -\sum_{a,a} - \sum_{b,b} + \sum_{c,c}$
   $\theta_4 = -\sum_{a,a} - \sum_{b,b} - \sum_{c,c}$.

   We see that θ_1 and θ_4 are in the same $Aut(J_{3,1})$-orbit (since $\theta_1 = -\theta_4$), also θ_2 and θ_3 are in the same $Aut(J_{3,1})$-orbit (since $\theta_2 = -\theta_3$). It remains to check that if θ_1 and θ_2 are in the same $Aut(J_{3,1})$-orbit or not. We claim that θ_1 and θ_2 are not in the same $Aut(J_{3,1})$-orbit. Let

$\phi \in Aut(J_{3,1})$ and $\lambda \in \mathbb{R}^*$ such that $\phi\theta_1 = \lambda\theta_2$. Then

$$\begin{aligned}
a_{11}^2 + a_{21}^2 + a_{31}^2 &= \lambda \qquad (5)\\
a_{12}^2 + a_{22}^2 + a_{32}^2 &= \lambda \qquad (6)\\
a_{13}^2 + a_{23}^2 + a_{33}^2 &= -\lambda \qquad (7)\\
a_{11}a_{12} + a_{21}a_{22} + a_{31}a_{32} &= 0\\
a_{11}a_{13} + a_{21}a_{23} + a_{31}a_{33} &= 0\\
a_{12}a_{13} + a_{22}a_{23} + a_{32}a_{33} &= 0
\end{aligned}$$

Equations $(5), (6)$ with (7) lead to $\lambda = 0$ and $\phi = 0$. So θ_1 and θ_2 are not in the same $Aut(J_{3,1})$-orbit, hence we get only two algebras

$J_{4,5}^{\alpha=\pm1}(\mathbb{R})$	$a^2 = d\ ,\ b^2 = d\ , c^2 = \alpha d.$

2. 1**- dimensional Central extension of** $J_{3,2}$

Here we get that $Z^2(J_{3,2}, K)$ is spanned by $\sum_{a,a}, \sum_{a,b}, \sum_{a,c}, \sum_{b,c}$ and $\sum_{c,c}$. Moreover, $\delta C^1(J_{3,2}, K)$ is spanned by $\sum_{a,a}$. Then $H^2(J_{3,2}, K)$ consists of $\theta := \alpha_1 \sum_{a,b} + \alpha_2 \sum_{a,c} + \alpha_3 \sum_{b,c} + \alpha_4 \sum_{c,c}$. The centre of $J_{3,2}, Z(J_{3,2})$, is spanned by b and c. Furthermore the automorphism group,$Aut(J_{3,2})$, consists of :

$$\phi = \begin{pmatrix} a_{11} & 0 & 0 \\ a_{21} & a_{11}^2 & a_{23} \\ a_{31} & 0 & a_{33} \end{pmatrix}\ ,\ \det\phi \neq 0.$$

The automorphism group acts as follows :

$$\begin{aligned}
\alpha_1 &\longrightarrow a_{11}^3\alpha_1 + a_{31}a_{11}^2\alpha_3\\
\alpha_2 &\longrightarrow a_{11}a_{23}\alpha_1 + a_{11}a_{33}\alpha_2 + (a_{21}a_{33} + a_{31}a_{23})\alpha_3 + a_{31}a_{33}\alpha_4\\
\alpha_3 &\longrightarrow a_{11}^2a_{33}\alpha_3\\
\alpha_4 &\longrightarrow 2a_{23}a_{33}\alpha_3 + a_{33}^2\alpha_4.
\end{aligned}$$

We distinguish two cases.

Case 1 : First suppose that $\alpha_3 \neq 0$, then we can divide to get $\alpha_3 = 1$. So we may assume that $\alpha_3 = 1$, choose $a_{11} = a_{33} = 1$, this leads to :

$$\begin{aligned}
\alpha_1 &\longrightarrow \alpha_1 + a_{31}\\
\alpha_2 &\longrightarrow a_{23}\alpha_1 + \alpha_2 + (a_{21} + a_{31}a_{23}) + a_{31}\alpha_4\\
\alpha_3 &\longrightarrow 1\\
\alpha_4 &\longrightarrow 2a_{23} + \alpha_4.
\end{aligned}$$

By taking $a_{31} = -\alpha_1$ and $a_{23} = -\frac{1}{2}\alpha_4$, we get $\alpha_1 \longrightarrow 0$ and $\alpha_4 \longrightarrow 0$. Then we can assume that $\alpha_1 = \alpha_4 = 0$ and $\alpha_3 = 1$. To conserve this we set $a_{31} = a_{23} = 0$ and $a_{11} = a_{33} = 1$, in this case we get

$$\begin{array}{lcl} \alpha_1 & \longrightarrow & 0 \\ \alpha_2 & \longrightarrow & \alpha_2 + a_{21} \\ \alpha_3 & \longrightarrow & 1 \\ \alpha_4 & \longrightarrow & 0. \end{array}$$

By taking $a_{21} = -\alpha_2$, then $\alpha_2 \longrightarrow 0$. Then we may assume that $\theta = \sum_{b,c}$, so we get the algebra :

$J_{4,6}(K)$	$a^2 = b\ ,\ b \circ c = d.$

$J_{4,6}(\mathbb{R})$	$a^2 = b\ ,\ b \circ c = d.$

Case 2 : If $\alpha_3 = 0$, this ledas to

$$\begin{array}{lcl} \alpha_1 & \longrightarrow & a_{11}^3\alpha_1 \\ \alpha_2 & \longrightarrow & a_{11}a_{23}\alpha_1 + a_{11}a_{33}\alpha_2 + a_{31}a_{33}\alpha_4 \\ \alpha_3 & \longrightarrow & 0 \\ \alpha_4 & \longrightarrow & a_{33}^2\alpha_4. \end{array}$$

In order to have $\theta^{\perp} \cap Z(J_{3,2}) = 0$, we need $\alpha_1 \neq 0$ and $(\alpha_2, \alpha_4) \neq (0,0)$. So after dividing we may assume $\alpha_1 = 1$. Choose $a_{11} = a_{33} = 1, a_{31} = 0$ and $a_{23} = -\alpha_2$, this leads to :

$$\begin{array}{lcl} \alpha_1 & \longrightarrow & 1 \\ \alpha_2 & \longrightarrow & 0 \\ \alpha_3 & \longrightarrow & 0 \\ \alpha_4 & \longrightarrow & \alpha_4 \neq 0. \end{array}$$

then we get the cocycle $\theta = \sum_{a,b} + \alpha_4 \sum_{c,c}$. Now the problem is to describe exactly when there exist $\phi \in Aut(J_{3,2})$ with $\phi(\sum_{a,b} + \alpha_4 \sum_{c,c}) = \lambda(\sum_{a,b} + \alpha_4^{'} \sum_{c,c})$ for some $\lambda \in K^*$. Then we get $\phi(\sum_{a,b} + \alpha_4 \sum_{c,c}) = \lambda(\sum_{a,b} + \alpha_4^{'} \sum_{c,c})$ if and only if there exist $a_{11}, a_{33} \in K^*$ such that $a_{33}^2\alpha_4 = a_{11}^3\alpha_4^{'}$. Then there is only one $Aut(J_{3,2})$-orbit, with representative $\theta = \sum_{a,b} + \sum_{c,c}$. So we get the algebra

$J_{4,7}(K)$	$a^2 = b\ ,\ a \circ b = d\ ,\ c^2 = d.$

$J_{4,7}(\mathbb{R})$	$a^2 = b\ ,\ a \circ b = d\ ,\ c^2 = d.$

Remarks :

- *For any* $\alpha, \beta \in K^*$ there exist $x, y \in K^*$ such that $x^3\alpha = y^2\beta$, by taking $x = \alpha\beta$ and $y = \alpha^2\beta$.
- Let $J^{\alpha}_{4,7} : a^2 = b\ ,\ a \circ b = d\ ,\ c^2 = \alpha d$. Now putting $a^{'} = \alpha a, b^{'} = \alpha^2 b, c^{'} = \gamma c, d^{'} = \alpha^3 d$ we get the same multiplication table, but the parameter has changed to 1.So we may assume $\alpha = 1$, and we get only one algebra $J_{4,7}$.
- Cocycles $\sum_{a,b} + \sum_{c,c}$ and $\sum_{b,c}$ not lie in the same $Aut(J_{3,2})$-orbit, $\phi\left(\sum_{a,b} + \sum_{c,c}\right) = a_{11}^3 \sum_{a,b} + (a_{11}a_{23} + a_{31}a_{33}) \sum_{a,c} + a_{33}^2 \sum_{c,c}$. Clearly $J_{4,6}$ and $J_{4,7}$ are not isomorphic, $J_{4,6}$ not associative but $J_{4,7}$ is a commutative nilpotent associative algebra.

3. **1- dimensional Central extension of** $J^{\alpha}_{3,3}, \alpha \in K^* \diagup (K^*)^2$

Here we get that $Z^2(J^{\alpha}_{3,3}, K)$ consists of :

$\theta := \alpha_1 \sum_{a,a} + \alpha_2 \sum_{b,b} + \alpha_3 \sum_{a,b} + \alpha_4 \sum_{a,c} + \alpha_5 \sum_{b,c}$.

Moreover, $\delta C^1(J^{\alpha}_{3,3}, K)$ is spanned by $\lambda \sum_{a,a} + \beta \sum_{b,b}$ for some $\lambda, \beta \in K$ such that $(\lambda, \beta) \neq (0, 0)$ and $\delta C^1(J^{\alpha}_{3,3}, K)$ is invarient under the action of $Aut(J^{\alpha}_{3,3})$. The automorphism group,$Aut(J^{\alpha}_{3,3})$, consists of :

$$\phi = \begin{pmatrix} a_{11} & a_{12} & 0 \\ a_{21} & a_{22} & 0 \\ a_{31} & a_{32} & a_{33} \end{pmatrix}$$

such that

$$\begin{aligned} a_{11}^2 + \alpha a_{21}^2 &= a_{33}\ ,\ a_{12}^2 + \alpha a_{22}^2 = a_{33} \\ a_{11}a_{12} + \alpha a_{21}a_{22} &= 0\ ,\ a_{33}\delta = a_{33}(a_{11}a_{22} - a_{21}a_{12}) \neq 0. \end{aligned}$$

Write $\phi\theta = \alpha_1' \sum_{a,a} + \alpha_2' \sum_{b,b} + \alpha_3' \sum_{a,b} + \alpha_4' \sum_{a,c} + \alpha_5' \sum_{b,c}$. Then

$$\begin{aligned} \alpha_1^{'} &= a_{11}^2\alpha_1 + a_{21}^2\alpha_2 + 2a_{11}a_{21}\alpha_2 + a_{11}a_{31}\alpha_4 + a_{21}a_{31}\alpha_5 \\ \alpha_2^{'} &= a_{12}^2\alpha_1 + a_{22}^2\alpha_2 + 2a_{11}a_{21}\alpha_2 + a_{12}a_{32}\alpha_4 + a_{22}a_{32}\alpha_5 \\ \alpha_3^{'} &= a_{11}a_{12}\alpha_1 + a_{22}a_{21}\alpha_2 + (a_{11}a_{22} + a_{21}a_{12})\alpha_3 \\ &\quad + (a_{11}a_{32} + a_{31}a_{12})\alpha_4 + (a_{21}a_{32} + a_{31}a_{22})\alpha_5 \\ \alpha_4^{'} &= \delta\left(a_{22}\alpha\alpha_3 - a_{12}\alpha_4\right) \\ \alpha_5^{'} &= a_{33}\left(a_{12}\alpha_3 + a_{22}\alpha_4\right). \end{aligned}$$

Now we need to determine λ and β, we have :

$\phi\left(\lambda \sum_{a,a} + \beta \sum_{b,b}\right) = (a_{11}^2\lambda + a_{21}^2\beta) \sum_{a,a} + (a_{12}^2\lambda + a_{22}^2\beta) \sum_{b,b} + a_{21}a_{22}\left(\beta - \alpha\lambda\right) \sum_{a,b}$.

It follows that $\delta C^1(J_{3,2},K)$ is invarient under the action of $Aut(J_{3,3}^{\alpha})$ if and only if $\beta=\alpha\lambda$, then $\delta C^1(J_{3,2},K)$ is spanned by $\sum_{a,a}+\alpha\sum_{b,b}$ where $\alpha\in K^{*}\diagup\left(K^{*}\right)^{2}$. Then $H^2(J_{3,3}^{\alpha},K)$ is consists of :

$\theta:=\alpha_1\sum_{a,a}+\alpha_2\sum_{b,b}+\alpha_3\sum_{a,b}+\alpha_4\sum_{a,c}+\alpha_5\sum_{b,c}$ modulo $\sum_{a,a}+\alpha\sum_{b,b}$.

The centre of $J_{3,3}^{\alpha}, Z(J_{3,3}^{\alpha})$, is spanned by c, so in order to have $\theta^{\perp}\cap Z(J_{3,3}^{\alpha})=0$ we need one of α_4,α_5 nonzero. Without loss of generality, we can assume that $\alpha_4=1$. Then

$\theta:=\alpha_1\sum_{a,a}+\alpha_2\sum_{b,b}+\alpha_3\sum_{a,b}+\sum_{a,c}+\alpha_5\sum_{b,c}$ modulo $\sum_{a,a}+\alpha\sum_{b,b}$.

Case $\alpha=1$:

We get

$$\begin{aligned}
\alpha_1^{'} &= a_{11}^2\alpha_1+a_{21}^2\alpha_2+2a_{11}a_{21}\alpha_2+a_{11}a_{31}+a_{21}a_{31}\alpha_5\\
\alpha_2^{'} &= a_{12}^2\alpha_1+a_{22}^2\alpha_2+2a_{11}a_{21}\alpha_2+a_{12}a_{32}+a_{22}a_{32}\alpha_5\\
\alpha_3^{'} &= a_{11}a_{12}\alpha_1+a_{22}a_{21}\alpha_2+(a_{11}a_{22}+a_{21}a_{12})\alpha_3\\
&\quad +a_{11}a_{32}+a_{31}a_{12}+(a_{21}a_{32}+a_{31}a_{22})\alpha_5\\
\alpha_4^{'} &= \delta\left(a_{22}-a_{12}\alpha_5\right)\\
\alpha_5^{'} &= a_{33}\left(a_{12}+a_{22}\alpha_5\right).
\end{aligned}$$

If $1+\alpha_5^2\neq 0$:

We can choose $a_{22}=\frac{1}{\delta\left(1+\alpha_5^2\right)}$ and $a_{12}=\frac{-\alpha_5}{\delta\left(1+\alpha_5^2\right)}$ to get $\alpha_4^{'}=1$ and $\alpha_5^{'}=0$. So may assume that $\alpha_4=1$ and $\alpha_5=0$, to fix θ, choose $a_{12}=a_{21}=0$ and $a_{11}=a_{22}=1$. It follows that

$$\begin{aligned}
\alpha_1^{'} &= \alpha_1+a_{31}\\
\alpha_2^{'} &= \alpha_2\\
\alpha_3^{'} &= \alpha_3+a_{32}\\
\alpha_4^{'} &= 1\\
\alpha_5^{'} &= 0.
\end{aligned}$$

Choose $a_{31}=\alpha_2-\alpha_1$ and $a_{32}=-\alpha_3$.

So we can assume that $\theta=\alpha_2\left(\sum_{a,a}+\sum_{b,b}\right)+\sum_{a,c}$ modulo $\sum_{a,a}+\sum_{b,b}$, then $\theta=\sum_{a,c}$. We get algebra

$J_{4,8}(K)$	$a^2=c\ ,\ b^2=c\ ,\ a\circ c=d.$

$J_{4,8}(\mathbb{R})$	$a^2=c\ ,\ b^2=c\ ,\ a\circ c=d.$

If $1+\alpha_5^2=0$:

Over algebraic closed field K and characteristic $K\neq 2$:

We have

$$\begin{array}{rcl}
\alpha_1^{'} & = & a_{11}^2\alpha_1+a_{21}^2\alpha_2+2a_{11}a_{21}\alpha_2+a_{11}a_{31}+a_{21}a_{31}\alpha_5 \\
\alpha_2^{'} & = & a_{12}^2\alpha_1+a_{22}^2\alpha_2+2a_{11}a_{21}\alpha_2+a_{12}a_{32}+a_{22}a_{32}\alpha_5 \\
\alpha_3^{'} & = & a_{11}a_{12}\alpha_1+a_{22}a_{21}\alpha_2+(a_{11}a_{22}+a_{21}a_{12})\alpha_3 \\
 & & +a_{11}a_{32}+a_{31}a_{12}+(a_{21}a_{32}+a_{31}a_{22})\alpha_5 \\
\alpha_4^{'} & = & \delta\left(a_{22}-a_{12}\alpha_5\right) \\
\alpha_5^{'} & = & a_{33}\left(a_{12}+a_{22}\alpha_5\right).
\end{array}$$

Choose $a_{12}=a_{21}=0, a_{11}=a_{22}=1, a_{31}=-\alpha_1$ and $a_{32}=\frac{-\alpha_2}{\alpha_5}$ (note that $\alpha_5\neq 0$ since $1+\alpha_5^2=0$). It follows that $\alpha_1^{'}=\alpha_2^{'}=0, \alpha_4^{'}=1, \alpha_5^{'}=\alpha_5$ and $1+\alpha_5^2=1+\alpha_5^{\prime 2}=0$.

So we assume that $\theta:=\alpha_3\sum_{a,b}+\sum_{a,c}+\alpha_5\sum_{b,c}$ such that $1+\alpha_5^2=0$. If $\alpha_3=0$ we get the algebra :

$$\boxed{a^2=c\ ,\ b^2=c\ ,\ a\circ c=d, b\circ c=\alpha_5 d, 1+\alpha_5^2=0.}$$

The change of basis $a\longrightarrow a, b\longrightarrow -\alpha_5 b, c\longrightarrow c$ and $d\longrightarrow d$, yields to

$J_{4,9}(K)$	$a^2=c\ ,\ b^2=-c\ ,\ a\circ c=d\ ,\ b\circ c=d\ .$

If $\alpha_3\neq 0$, by consider the previous change of the basis we get the algebra :

$$\boxed{a^2=c\ ,\ b^2=-c\ ,\ a\circ c=d\ , b\circ c=d\ ,\ a\circ b=\alpha_3 d.}$$

Now putting $a'=\alpha_3 a, b'=\alpha_3 b, c'=\alpha_3^2 c, d'=\alpha_3^3 d$ we get the same multiplication table, but the parameter has changed to 1. So we may assume $\alpha_3=1$, and we get only algebra

$J_{4,10}(K)$	$a^2=c\ ,\ b^2=-c\ ,\ a\circ c=d\ ,\ b\circ c=d\ ,\ a\circ b=d.$

Over real field $\mathbb{R}$:

There is no $\alpha_5\in\mathbb{R}$ such that $1+\alpha_5^2=0$.

We are completed 1- **d**imensional Central extension of $J_{3,3}^{\alpha}$ $\alpha\in K^*\diagup\left(K^*\right)^2$over algebraic closed field K and characteristic $K\neq 2$ since $K^*\diagup\left(K^*\right)^2=\{1\}$. To complete classification over $\mathbb{R}$, it remains to consider case $\alpha=-1$.

Case $\alpha = -1$ **:**

We get :

$$\begin{aligned}
\alpha_1^{'} &= a_{11}^2\alpha_1 + a_{21}^2\alpha_2 + 2a_{11}a_{21}\alpha_2 + a_{11}a_{31} + a_{21}a_{31}\alpha_5 \\
\alpha_2^{'} &= a_{12}^2\alpha_1 + a_{22}^2\alpha_2 + 2a_{11}a_{21}\alpha_2 + a_{12}a_{32} + a_{22}a_{32}\alpha_5 \\
\alpha_3^{'} &= a_{11}a_{12}\alpha_1 + a_{22}a_{21}\alpha_2 + (a_{11}a_{22} + a_{21}a_{12})\alpha_3 \\
&\quad + a_{11}a_{32} + a_{31}a_{12} + (a_{21}a_{32} + a_{31}a_{22})\alpha_5 \\
\alpha_4^{'} &= -\delta\left(a_{22} + a_{12}\alpha_5\right) \\
\alpha_5^{'} &= a_{33}\left(a_{12} + a_{22}\alpha_5\right).
\end{aligned}$$

As we do before

If $1-\alpha_5^2 \neq 0$:

We can choose $a_{22} = \frac{-1}{\delta\left(1-\alpha_5^2\right)}$ and $a_{12} = \frac{\alpha 5}{\delta\left(1-\alpha_5^2\right)}$ to get $\alpha_4^{'} = 1$ and $\alpha_5^{'} = 0$. So may assume that $\alpha_4 = 1$ and $\alpha_5 = 0$, to fix θ, choose $a_{12} = a_{21} = 0$ and $a_{11} = a_{22} = 1$. It follows that

$$\begin{aligned}
\alpha_1^{'} &= \alpha_1 + a_{31} \\
\alpha_2^{'} &= \alpha_2 \\
\alpha_3^{'} &= \alpha_3 + a_{32} \\
\alpha_4^{'} &= 1 \\
\alpha_5^{'} &= 0.
\end{aligned}$$

Choose $a_{31} = -\alpha_2 - \alpha_1$ and $a_{32} = -\alpha_3$.

So we can assume that $\theta = -\alpha_2\left(\sum_{a,a} - \sum_{b,b}\right) + \sum_{a,c}$ modulo $\sum_{a,a} - \sum_{b,b}$, then $\theta = \sum_{a,c}$. We get algebra

$J_{4,9}(\mathbb{R})$	$a^2 = c\ ,\ b^2 = -c\ ,\ a \circ c = d.$

If $1-\alpha_5^2 = 0$:

We have

$$\begin{aligned}
\alpha_1^{'} &= a_{11}^2\alpha_1 + a_{21}^2\alpha_2 + 2a_{11}a_{21}\alpha_2 + a_{11}a_{31} + a_{21}a_{31}\alpha_5 \\
\alpha_2^{'} &= a_{12}^2\alpha_1 + a_{22}^2\alpha_2 + 2a_{11}a_{21}\alpha_2 + a_{12}a_{32} + a_{22}a_{32}\alpha_5 \\
\alpha_3^{'} &= a_{11}a_{12}\alpha_1 + a_{22}a_{21}\alpha_2 + (a_{11}a_{22} + a_{21}a_{12})\alpha_3 \\
&\quad + a_{11}a_{32} + a_{31}a_{12} + (a_{21}a_{32} + a_{31}a_{22})\alpha_5 \\
\alpha_4^{'} &= -\delta\left(a_{22} + a_{12}\alpha_5\right) \\
\alpha_5^{'} &= a_{33}\left(a_{12} + a_{22}\alpha_5\right).
\end{aligned}$$

Choose $a_{12} = a_{21} = 0, a_{11} = a_{22} = 1, a_{31} = -\alpha_1$ and $a_{32} = \frac{-\alpha_2}{\alpha_5}$ (note that $\alpha_5 \neq 0$ since $1 - \alpha_5^2 = 0$). It follows that $\alpha_1' = \alpha_2' = 0, \alpha_4' = 1, \alpha_5' = \alpha_5$ and $1 - \alpha_5^2 = 1 - \alpha_5'^2 = 0$.

So we assume that $\theta := \alpha_3 \sum_{a,b} + \sum_{a,c} + \alpha_5 \sum_{b,c}$ such that $1 - \alpha_5^2 = 0$. If $\alpha_3 = 0$ we get the algebra :

$$\boxed{a^2 = c \ , \ b^2 = -c \ , \ a \circ c = d, b \circ c = \alpha_5 d, 1 - \alpha_5^2 = 0.}$$

The change of basis $a \longrightarrow a, b \longrightarrow \alpha_5 b, c \longrightarrow c$ and $d \longrightarrow d$, yields to

$J_{4,10}(\mathbb{R})$	$a^2 = c \ , \ b^2 = -c \ , \ a \circ c = d \ , \ b \circ c = d$.

If $\alpha_3 \neq 0$, by consider the previous change of the basis we get the algebra :

$$\boxed{a^2 = c \ , \ b^2 = -c \ , \ a \circ c = d \ , b \circ c = d \ , \ a \circ b = \alpha_3 d.}$$

Now putting $a' = \alpha_3 a, b' = \alpha_3 b, c' = \alpha_3^2 c, d' = \alpha_3^3 d$ we get the same multiplication table, but the parameter has changed to 1. So we may assume $\alpha_3 = 1$, and we get only algebra

$J_{4,11}(\mathbb{R})$	$a^2 = c \ , \ b^2 = -c \ , \ a \circ c = d \ , \ b \circ c = d \ , \ a \circ b = d.$

Remark :

One can consider the algebra $J : a \circ b = c$ instead of $J_{3,3}^{\alpha}$ over algebaic closed field K and $ch(K) \neq 2$, by change of basis $a \longrightarrow a + \sigma b, b \longrightarrow a - \sigma b, c \longrightarrow 2c$ and $\sigma^2 + 1 = 0$. Also for $J_{3,3}^{\alpha}, \alpha = -1$ in the case of the Real field $\mathbb{R}$, by change of basis $a \longrightarrow a + b, b \longrightarrow a - b, c \longrightarrow 2c$. Calculations by this consideration may by some simple. Here we get that $H^2(J, K)$ consists of $\theta = \alpha_1 \sum_{a,a} + \alpha_2 \sum_{b,b} + \alpha_3 \sum_{a,c} + \alpha_4 \sum_{b,c}$, and the automorphism group consists of

$$\phi = \begin{pmatrix} a_{11} & a_{12} & 0 \\ a_{21} & a_{22} & 0 \\ a_{31} & a_{32} & a_{11}a_{22} + a_{21}a_{12} \end{pmatrix}, a_{11}^2 a_{22}^2 - a_{12}^2 a_{21}^2 \neq 0 \text{ and } \ a_{11}a_{21} = a_{12}a_{22} = 0$$

Then $\phi\theta = \alpha_1' \sum_{a,a} + \alpha_2' \sum_{b,b} + \alpha_3' \sum_{a,c} + \alpha_4' \sum_{b,c}$

$$\begin{aligned} \alpha_1' &= a_{11}^2 \alpha_1 + 2a_{11}a_{31}\alpha_3 \\ \alpha_2' &= a_{22}^2 \alpha_2 + 2a_{22}a_{32}\alpha_4 \\ \alpha_3' &= (a_{11}a_{22} + a_{21}a_{12})(a_{11}\alpha_3 + a_{21}\alpha_4) \\ \alpha_4' &= (a_{11}a_{22} + a_{21}a_{12})(a_{12}\alpha_3 + a_{22}\alpha_4) \end{aligned}$$

We find that there exist only three orbits, $\theta_1 = \sum_{a,c}, \theta_2 = \sum_{a,c} + \sum_{b,b}$ and $\theta_3 = \sum_{a,c} + \sum_{b,c}$, which be the same number of orbits we get for these algebras.

4. 1- **dimensional Central extension of** $J_{3,4}$

Here $Z^2(J_{3,4}, K)$ is spanned by $\sum_{a,a}, \sum_{a,b}$ and $\sum_{a,c} + \sum_{b,b}$. Moreover, $\delta C^1(J_{3,4}, K)$ is spanned by $\sum_{a,a}$ and $\sum_{a,b}$. Then $H^2(J_{3,4}, K)$ is spanned by $\theta := \sum_{a,c} + \sum_{b,b}$, so we get only one algebra

$J_{4,11}(K)$	$a^2 = b \ , \ a \circ b = c \ , \ a \circ c = d \ , \ b^2 = d.$

$J_{4,12}(\mathbb{R})$	$a^2 = b \ , \ a \circ b = c \ , \ a \circ c = d \ , \ b^2 = d.$

5. 2- **dimensional Central extension of** $J_{2,1}$

Here $H^2(J_{2,1}, K)$ is spanned by $\sum_{a,a}, \sum_{a,b}$ and $\sum_{b,b}$, then $G_2(H^2(J_{2,1}, K))$ consists of :

$\theta := \alpha_1 \sum_{a,a} \wedge \sum_{a,b} + \alpha_2 \sum_{a,a} \wedge \sum_{b,b} + \alpha_3 \sum_{a,b} \wedge \sum_{b,b}$.

The autommorphism group,$Aut(J_{2,1})$, consists of :

$$\phi = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}, \det \phi \neq 0.$$

Now we need to find representatives of $U_2(J_{2,1}) \diagup Aut(J_{2,1})$, write

$\phi\theta := \alpha_1' \sum_{a,a} \wedge \sum_{a,b} + \alpha_2' \sum_{a,a} \wedge \sum_{b,b} + \alpha_3' \sum_{a,b} \wedge \sum_{b,b}$.

Then

$$\begin{aligned} \alpha_1' &= \left(a_{11}^2\alpha_1 + a_{11}a_{21}\alpha_2 + a_{21}^2\alpha_3\right) \det \phi \\ \alpha_2' &= \left(2a_{11}a_{12}\alpha_1 + (a_{11}a_{22} + a_{12}a_{21})\alpha_2 + 2a_{21}a_{22}\alpha_3\right) \det \phi \\ \alpha_3' &= \left(a_{12}^2\alpha_1 + a_{12}a_{22}\alpha_2 + a_{22}^2\alpha_3\right) \det \phi. \end{aligned}$$

Without loss of generality, we can assume that $\alpha_1 = 1$ since $(\alpha_1, \alpha_2, \alpha_3) \neq (0, 0, 0)$. Choose $a_{11} = a_{22} = 1, a_{21} = 0, a_{12} = \frac{-\alpha_2}{2}$ to get $\alpha_1' = 1$ and $\alpha_2' = 0$.

So we may assume $\theta = \sum_{a,a} \wedge \sum_{a,b} + \alpha_3 \sum_{a,b} \wedge \sum_{b,b}$.

If $\alpha_3 = 0$:

Then we get the cocycle $\theta_1 = \sum_{a,a} \wedge \sum_{a,b}$.

If $\alpha_3 \neq 0$:

$$\begin{aligned} \alpha_1' &= \left(a_{11}^2\alpha_1 + a_{21}^2\alpha_3\right) \det \phi \\ \alpha_2' &= \left(2a_{11}a_{12}\alpha_1 + 2a_{21}a_{22}\alpha_3\right) \det \phi \\ \alpha_3' &= \left(a_{12}^2\alpha_1 + a_{22}^2\alpha_3\right) \det \phi. \end{aligned}$$

To fix $\alpha'_2 = 0$ choose $a_{21} = a_{12} = 0$, then

$$\begin{aligned}\alpha'_1 &= \det\phi\left(a_{11}^2\right)\\ \alpha'_2 &= 0\\ \alpha'_3 &= \det\phi\left(a_{22}^2\alpha_3\right).\end{aligned}$$

Over algebraic closed field K and characteristic $K \neq 2$:

In this case every element is a square, then we choose $a_{11} = 1, a_{22} = \frac{1}{\sqrt{-\alpha_3}}$ and then dividing by $\det\phi$. We get $\theta_2 = \sum_{a,a} \wedge \sum_{a,b} - \sum_{a,b} \wedge \sum_{b,b} = \left(\sum_{a,a} + \sum_{b,b}\right) \wedge \sum_{a,b}$. We claim that θ_1 and θ_2 are not in the same orbit. Let $\phi \in Aut(J_{2,1})$ and $\lambda \in K^*$ such that $\phi\ \theta_1 = \lambda\theta_2$,then

$$\begin{aligned}\lambda &= \left(a_{11}^2\right)\det\phi\\ 0 &= (2a_{11}a_{12})\det\phi \qquad (8)\\ -\lambda &= \left(a_{12}^2\right)\det\phi.\end{aligned}$$

Equation (8) ensures that $\lambda = 0$. So we get only two $Aut(J_{2,1})$-orbits over K, yielding the two algebras

$J_{4,12}(K)$	$a^2 = c\ ,\ a \circ b = d\ .$

$J_{4,13}(K)$	$a^2 = c\ ,\ b^2 = c\ ,\ a \circ b = d\ .$

Over real field $\mathbb{R}$:

If $\alpha_3 < 0$, choose $a_{11} = 1, a_{22} = \frac{1}{\sqrt{-\alpha_3}}$ and then dividing by $\det\phi$.We get :

$\theta_2 = \sum_{a,a} \wedge \sum_{a,b} - \sum_{a,b} \wedge \sum_{b,b} = \left(\sum_{a,a} + \sum_{b,b}\right) \wedge \sum_{a,b}$.

If $\alpha_3 > 0$, choose $a_{11} = 1, a_{22} = \frac{1}{\sqrt{\alpha_3}}$ and then dividing by $\det\phi$. We get :

$\theta_3 = \sum_{a,a} \wedge \sum_{a,b} + \sum_{a,b} \wedge \sum_{b,b} = \left(\sum_{a,a} - \sum_{b,b}\right) \wedge \sum_{a,b}$.

Clearly θ_1 is not conjugate to any of θ_2 and θ_3. It remains to check if θ_2 and θ_3 are in the same $Aut(J_{2,1})$-orbit or not. Let $\phi \in Aut(J_{2,1})$ and $\lambda \in \mathbb{R}^*$ such that $\phi\ \theta_3 = \lambda\theta_2$,then

$$\begin{aligned}\lambda &= \left(a_{11}^2 + a_{21}^2\right)\det\phi \qquad (9)\\ 0 &= (2a_{11}a_{12} + 2a_{21}a_{22})\det\phi\\ -\lambda &= \left(a_{12}^2 + a_{22}^2\right)\det\phi. \qquad (10)\end{aligned}$$

Equations (9) and (10) Lead to $\lambda = 0$ and $\phi = 0$. So we get only three $Aut(J_{2,1})$-orbits over $\mathbb{R}$, yielding the three algebras

$J_{4,13}(\mathbb{R})$	$a^2 = c \ , \ a \circ b = d \ .$

$J_{4,14}(\mathbb{R})$	$a^2 = c \ , \ b^2 = c \ , a \circ b = d \ .$

$J_{4,15}(\mathbb{R})$	$a^2 = c \ , \ b^2 = -c \ , a \circ b = d \ .$

Summarize our results in the following theorems :

Theorem 37 *Up to isomorphism there exist* 13 *nilpotent Jordan algebras of dimension* 4 *over algebraic closed field* K *and* $ch(K) \neq 2$ *which are isomorphic to one of the following pairwise non-isomorphic nilpotent Jordan algebras :*

Nilpotent Jordan algebras with Centeral component
$J_{4,1} = J_{3,1} \oplus J_{1,1}$. :*All multiplications are zero.*
$J_{4,2} = J_{3,2} \oplus J_{1,1}. : a^2 = b.$
$J_{4,3} = J_{3,3} \oplus J_{1,1}. : a^2 = c \ , b^2 = c.$
$J_{4,4} = J_{3,4} \oplus J_{1,1}. : a^2 = b \ , a \circ b = c.$
Nilpotent Jordan algebras without Centeral component
$J_{4,5} : a^2 = d \ , b^2 = d \ , c^2 = d..$
$J_{4,6} : a^2 = b \ , \ b \circ c = d.$
$J_{4,7} : a^2 = b \ , \ a \circ b = d \ , \ c^2 = d.$
$J_{4,8} : a^2 = c \ , \ b^2 = c \ , \ a \circ c = d.$
$J_{4,9} : a^2 = c \ , \ b^2 = -c \ , \ a \circ c = d \ , b \circ c = d \ .$
$J_{4,10} : a^2 = c \ , \ b^2 = -c \ , \ a \circ c = d \ , b \circ c = d \ , a \circ b = d.$
$J_{4,11} : a^2 = b \ , \ a \circ b = c \ , a \circ c = d \ , b^2 = d.$
$J_{4,12} : a^2 = c \ , \ a \circ b = d \ .$
$J_{4,13} : a^2 = c \ , \ b^2 = c \ , a \circ b = d \ .$

We see that all nilpotent Jordan algebras of dimension 4 over algebraic closed field K and $ch(K) \neq 2$ are associative except $J_{4,6}, J_{4,8}, J_{4,9}$ and $J_{4,10}$.

Theorem 38 *Up to isomorphism there exist* 9 *commutative nilpotent associative algebras of dimension* 4 *over algebraic closed field* K *and* $ch(K) \neq 2$ *which are isomorphic to one of the following pairwise non-isomorphic com-*

mutative nilpotent associative algebras :

Nilpotent Jordan algebras with Centeral component
$J_{4,1} = J_{3,1} \oplus J_{1,1}.$:*All multiplications are zero.*
$J_{4,2} = J_{3,2} \oplus J_{1,1}.$: $a^2 = b.$
$J_{4,3} = J_{3,3} \oplus J_{1,1}.$: $a^2 = c$, $b^2 = c.$
$J_{4,4} = J_{3,4} \oplus J_{1,1}.$: $a^2 = b$, $a \circ b = c.$
Nilpotent Jordan algebras without Centeral component
$J_{4,5}$: $a^2 = d$, $b^2 = d$,$c^2 = d..$
$J_{4,7}$: $a^2 = b$, $a \circ b = d$, $c^2 = d.$
$J_{4,11}$: $a^2 = b$, $a \circ b = c$, $a \circ c = d$, $b^2 = d.$
$J_{4,12}$: $a^2 = c$, $a \circ b = d$.
$J_{4,13}$: $a^2 = c$, $b^2 = c$, $a \circ b = d$.

Theorem 39 *Up to isomorphism there exist* 17 *nilpotent Jordan algebras of dimension* 4 *over* $\mathbb{R}$ *which are isomorphic to one of the following pairwise non-isomorphic nilpotent Jordan algebras :*

Nilpotent Jordan algebras with Centeral component
$J_{4,1} = J_{3,1} \oplus J_{1,1}.$:*All multiplications are zero.*
$J_{4,2} = J_{3,2} \oplus J_{1,1}.$: $a^2 = b.$
$J_{4,3}^{\alpha=\pm1} = J_{3,3}^{\alpha=\pm1} \oplus J_{1,1}.$: $a^2 = c$, $b^2 = \alpha c$. $(\alpha = \pm1)$
$J_{4,4} = J_{3,4} \oplus J_{1,1}.$: $a^2 = b$, $a \circ b = c.$
Nilpotent Jordan algebras without Centeral component
$J_{4,5}^{\alpha=\pm1}$: $a^2 = d$, $b^2 = d$,$c^2 = \alpha d$. $(\alpha = \pm1)$
$J_{4,6}$: $a^2 = b$, $b \circ c = d.$
$J_{4,7}$: $a^2 = b$, $a \circ b = d$, $c^2 = d.$
$J_{4,8}^{\alpha=\pm1}$: $a^2 = c$, $b^2 = \alpha c$, $a \circ c = d.$ $(\alpha = \pm1)$
$J_{4,9}$: $a^2 = c$, $b^2 = -c$, $a \circ c = d$, $b \circ c = d$.
$J_{4,10}$: $a^2 = c$, $b^2 = -c$, $a \circ c = d$, $b \circ c = d$, $a \circ b = d.$
$J_{4,11}$: $a^2 = b$, $a \circ b = c$, $a \circ c = d$, $b^2 = d.$
$J_{4,12}$: $a^2 = c$, $a \circ b = d$.
$J_{4,13}^{,\alpha=\pm1.}$: $a^2 = c$, $b^2 = \alpha c$, $a \circ b = d$. $(\alpha = \pm1)$

Also, all nilpotent Jordan algebras of dimension 4 over $\mathbb{R}$ are associative except $J_{4,6}, J_{4,8}^{\alpha=\pm1}, J_{4,9}$ and $J_{4,10}$.

Theorem 40 *Up to isomorphism there exist* 12 *commutative nilpotent associative algebras of dimension* 4 *over* $\mathbb{R}$ *which are isomorphic to one of the* *following pairwise non-isomorphic commutative nilpotent associative algebras :*

Nilpotent Jordan algebras with Centeral component
$J_{4,1} = J_{3,1} \oplus J_{1,1}.$:*All multiplications are zero.*
$J_{4,2} = J_{3,2} \oplus J_{1,1}.$: $a^2 = b.$
$J_{4,3}^{\alpha=\pm1} = J_{3,3}^{\alpha=\pm1} \oplus J_{1,1}.$: $a^2 = c$, $b^2 = \alpha c$. $(\alpha = \pm 1)$
$J_{4,4} = J_{3,4} \oplus J_{1,1}.$: $a^2 = b$, $a \circ b = c.$
Nilpotent Jordan algebras without Centeral component
$J_{4,5}^{\alpha=\pm1}$: $a^2 = d$, $b^2 = d$, $c^2 = \alpha d$. $(\alpha = \pm 1)$
$J_{4,7}$: $a^2 = b$, $a \circ b = d$, $c^2 = d.$
$J_{4,11}$: $a^2 = b$, $a \circ b = c$, $a \circ c = d$, $b^2 = d.$
$J_{4,12}$: $a^2 = c$, $a \circ b = d$.
$J_{4,13}^{,\alpha=\pm1.}$: $a^2 = c$, $b^2 = \alpha c$, $a \circ b = d$. $(\alpha = \pm 1)$

References

[1] ST. Skjelbred and T. Sund. On the classification of nilpotent Lie algebras. Technical report Mathematisk Institutt, Universitetet i Oslo, (1977).

[2] Gong M. P. Classification of Nilpotent Lie Algebras of Dimension 7. PhD thesis, University of Waterloo, Canada, 1998.

[3] R.E. Beck, B. Kolman, Construction of nilpotent Lie algebras over arbitrary fields, in: Paul S.Wang (Ed.), Proceedings of the 1981 ACM Symposium on Symbolic and Algebraic Computation, ACM, New York, 1981, pp. 169–174.

[4] de Graaf W. A. Classification of 6-dimensional nilpotent Lie algebras over fields of characteristic not 2, J. Algebra. 309(2)(2007), 640-653.

[5] D. Cox, J. Little, D. O'Shea, Ideals, Varieties and Algorithms: An Introduction to Computational Algebraic Geometry and Commutative Algebra, Springer, New York, 1992.

[6] A. A. Albert. A structure theory for Jordan algebras, Ann.Math. 48 (1946),546-567.

[7] W. A. de Graaf. Classification of nilpotent associative algebras of small dimension. arXiv:1009.5339v1 [math.RA], 2010.

[8] O. Timothy O'Meara. Introduction to quadratic forms. Classics in Mathematics. Springer Verlag, Berlin, 2000. Reprint of the 1973 edition.

[9] Nathan Jacobson. Lectures in abstract algebra. Springer-Verlag, New York, 1975. Volume II: Linear algebra, Reprint of the 1953 edition [Van Nostrand, Toronto, Ont.], Graduate Texts in Mathematics, No. 31.

[10] Milnor, J.; Husemoller, D. Symmetric Bilinear Forms. Springer-Verlag New York Heidelberg Berlin 1973,ISBN 0-387-06009-X .

[11] Jean-Pierre Serre. A Course in Arithmetic, Springer-Verlag, 1973.

[12] Schafer, Richard D. (1995) [1966], An Introduction to Nonassociative Algebras, Dover, ISBN 0-486-68813-5

[13] E. A. Dekerf , A. P. E Ten Kroode. Lie Algebras, Finite and Infinite Dimensional Lie Algebras and Applications in Physics part 2, Elsevier, ISBN 0444828362 .

[14] Karlheinz Spindler. Abstract Algebra With Applications, Volumes I, Marcel Dekker 1994. ISBN 0-8247-9144-4